

Review

Open Access

## Reproductive toxicity of seafood contaminants: Prospective comparisons of Swedish east and west coast fishermen's families

Anna Axmon\*, Lars Rylander and Anna Rignell-Hydbom

Address: Division of Occupational and Environmental Medicine, Department of Laboratory Medicine, Lund University, University Hospital, SE-221 85 Lund, Sweden

Email: Anna Axmon\* - [anna.axmon@med.lu.se](mailto:anna.axmon@med.lu.se); Lars Rylander - [lars.rylander@med.lu.se](mailto:lars.rylander@med.lu.se); Anna Rignell-Hydbom - [anna.rignell-hydbom@med.lu.se](mailto:anna.rignell-hydbom@med.lu.se)

\* Corresponding author

Published: 28 May 2008

Received: 10 October 2007

*Environmental Health* 2008, **7**:20 doi:10.1186/1476-069X-7-20

Accepted: 28 May 2008

This article is available from: <http://www.ehjournal.net/content/7/1/20>

© 2008 Axmon et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

Cohorts comprising fishermen's families on the east coast of Sweden have been found to have a high consumption of contaminated fish as well as high body burdens of persistent organochlorine pollutants (POPs). Their west coast correspondents are socio-economically similar, but with considerably lower POP exposure since the fish caught on the west coast is far less contaminated. The rationale for this was that the cohorts residing on the east coast of Sweden have been found to have a high consumption of contaminated fish as well as high body burdens of POPs, whereas their west coast correspondents are socio-economically similar, but with considerably lower POP exposure since the fish caught on the west coast is far less contaminated. Among the reproductive outcomes investigated are included both male and female parameters, as well as couple fertility and effects on the fetus. A range of exposure measures, including both questionnaire assessments of fish consumption and biomarkers, have been used.

The most consistent findings of the studies are those related to the fetus, where a decreased birth weight was found across all measures of exposure, which is in agreement with studies from other populations. Some markers for male reproduction function, i.e. sperm motility, sperm chromatin integrity, and Y:X chromosome ratio, were associated with POP exposure, whereas others, such as sperm concentration and semen volume, were not. With respect to couple fertility and female reproductive parameters, no support was given for associations with POP exposure. Although some associations may have been affected by beneficial effects of essential nutrients in seafood, the overall findings are meaningful in the context of reproductive toxicity and support the usefulness of the epidemiological design.

### Introduction

Concerns regarding health effects of persistent organochlorine pollutants (POPs) were first expressed in the 1960's when exposed animals were found to be affected, and exhibiting thinning eggshells, morphologic abnormalities, and impaired viability of offspring [1]. Since

then, the topic has been thoroughly investigated in experimental animal studies as well as epidemiological animal and human studies. Several POPs are endocrine disruptors, *i.e.* chemicals with sex steroid mimicking effects which may cause developmental and reproductive abnormalities in humans and animals [2-4].

The term "persistent organochlorine pollutants" (POPs) is often used to include several compounds, such as polychlorinated biphenyls (PCBs), dioxins and organochlorine insecticides such as 1,1,1-trichloro-2,2-bis(p-chlorophenyl)ethane (DDT). Although they have had different applications in the industry, POPs have several traits in common: They are lipophilic and poorly metabolized, and they bioaccumulate. While they are not very soluble in water, they have spread widely through the atmosphere, and are found in parts of the world where they were never used [5]. Moreover, even after their use has been banned, POPs are still found in human and animal tissue and in body fluids [6-10].

The most common exposure route for the majority of individuals is through the consumption of contaminated foods. In common, for some of the most highly exposed populations, such as Inuits from Greenland and individuals from Faroe Islands, is that the main exposure source is from the seafood [9,11]. In Sweden, the major exposure source is consumption of fatty fish from the Baltic Sea, off the east coast of the country. As a consequence, plasma levels of several of the compounds were higher in men with a high intake of such fish than in those who consumed only moderate amounts, and the levels were higher in those who ate moderate amounts of fish than in those who ate none [12].

It has been found that both professional east coast fishermen and their wives have a higher intake of Baltic Sea fish than referents from the same area [13,14]. Similarly, on the west coast of Sweden, fishermen and their wives have a higher consumption of locally caught fish than referents from the same area. The average amount of locally caught fish consumed by fishermen and their wives on the east and west coast, respectively, has been found to be similar. However, the fish caught off the west coast of Sweden has for the past decades been much less contaminated than that caught from the Baltic Sea [15]. Thus, east coast fishermen have been found to have higher levels of POPs than both west coast fishermen and referents [13,16]. Consequently, the east coast fishermen and their wives constitute a suitable exposed group for investigating health effects of adult dietary exposure to POPs, whereas their counterparts on the west coast is an appropriate referent group. The socio-economic similarities between these cohorts in combination with a broad exposure range of POPs, gave very good opportunity to investigate the hypothesized association between POP exposure and reproductive health. Moreover, east coast fishermen's sisters have been found to be more likely to have grown up in a fishing village and/or fisherman's family [17], suggesting that they may be a fitting group when investigating health effects due to a high dietary exposure to POPs during childhood and adolescence, or even in utero. There-

fore, the cohorts of fishermen, their wives and sisters, from the two coastal areas of Sweden (The Swedish Fishermen's Families Cohorts) have over the last two decades been used to study a variety of health effects.

#### **Study populations**

In 1988, two cohorts of fishermen from the east (Baltic Sea) and west (Kattegatt and Skagerrak) coasts of Sweden were established [13,18]. Cohort affiliation was determined by the particular organization the fisherman was associated with. All in all, 2883 east coast fishermen and 8477 west coast fishermen were included in the cohorts. However, a subgroup of 96 east and 99 west coast fishermen also participated in a second study, in which both semen and blood samples were collected [16,19-21].

The fishermen cohorts were linked to the national Swedish Population Register and to registers at the local parish offices [22]. Thus, 2175 east and 7062 west coast women who were, or had been, married to a man in one of the original fishermen cohorts were identified. Cohort affiliation was determined by that of the fisherman to whom the woman was associated. Subgroups of these women have supplied blood samples [23] and information on time to pregnancy (TTP) [24]. Moreover, the entire cohorts of fishermen's wives were linked to the Swedish Medical Birth Registry (MBR) [14], thus identifying 1501 infants born to the east coast women during 1973 through 1991, with the corresponding number for the west coast being 3553. The children were assigned the same cohort affiliation as their mothers.

In 1996, the cohorts of fishermen were linked to the National Swedish Population Register in order to identify sisters (at least one parent in common) to these men [17]. Women who were included in the fishermen's wives cohort were excluded. The cohort affiliation was determined by that of the fisherman to whom the woman was associated. The cohorts of sisters were then linked to the MBR, and 1719 east and 1537 west coast infants were identified. Also among the fishermen's sisters a study regarding TTP was performed, including 1241 east and 2023 west coast women [25]. Of these, 165 women also provided blood samples [26].

#### **Measures of exposure**

In the studies performed on the Swedish Fishermen's Families Cohorts, several measures of exposure have been used (c.f. Table 1). In those studies which rely on register data and/or questionnaires and telephone interviews, the main measure of exposure has been *cohort affiliation*, using the east coast cohorts as exposed cohorts and the west coast cohorts as referent cohorts. Moreover, within the east coast cohorts, *growing up in a fishing village and/or fisherman's family* has been used as a proxy measure of expo-

**Table 1: Reproductive outcomes investigated in the Swedish Fishermen's Families Cohorts.**

Unit of observation	Outcome <sup>1</sup>	Exposure				p,p'-DDE <sup>4</sup>
		Cohort affiliation	Childhood exposure <sup>2</sup>	Fish consumption	CB-153 <sup>3</sup>	
Man	Sperm motility	-	-	-	↓ [16]	↓ [16]
	Sperm concentration	-	-	-	0 [16]	0 [16]
	Semen volume	-	-	-	0 [16]	0 [16]
	Total sperm count	-	-	-	0 [16]	0 [16]
	Sperm chromatin integrity	-	-	-	↓ [20]	(↓) [20]
	Y:X chromosome ratio	-	-	-	↓ [21]	↓ [21]
	Seminal levels of markers of epididymal and accessory sex gland function	-	-	-	0 [19]	0 [19]
Woman	Hormone parameters	-	-	-	0 [16]	0 [16]
	Age at menarche	0 [38]	↓ [38]	-	-	-
	Menstrual cycle length	↓ [91]	0 [91]	(↓) [91]	0 [91]	-
Couple	Miscarriages and stillbirths	NC [17, 25, 40]	0 [25, 40]	0 [25]	NC [26, 40]	-
	Time to pregnancy	NC [24, 25]	0 [24, 25]	0 [24, 25]	0 [26, 33]	-
Infant/child	Birth weight	↓ [14, 17]	↓ [41]	↓ [41]	↓ [31]	-
	Birth length	0 [14, 17]	-	-	-	-
	Head circumference	↓ [14, 17]	-	-	-	-
	Gender ratio	NC [14, 17]	-	-	-	-
	Malformations	0 [17, 43]	-	-	-	-

<sup>1</sup> Arrows indicate the effect associated with each exposure, with parenthesis for those effects not found to be of a noteworthy size but not statistically significant. A "0" means that no effect was found, and "NC" signifies non conclusive results.

<sup>2</sup> Growing up in a fisherman's family and/or fishing village

<sup>3</sup> 2,2',4,4',5,5'-hexachlorobiphenyl

<sup>4</sup> 1,1-dichloro-2,2-bis (p-chlorophenyl)-ethylene

sure during childhood and adolescence. In one of the earlier studies on the fishermen's wives, it was found that recall of fish consumption was poor [23]. Thus, in subsequent studies, *current fish consumption* was used as a proxy measure of exposure during the time of interest for each specific outcome (e.g. pregnancy attempt or child birth).

With respect to the majority of the reproductive outcomes investigated, *biomarkers* have also been used as measures of exposure. The main biomarker has been 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153). This congener has been found to correlate strongly with the total PCB concentration, as well as the total PCB-derived TEQ and the total POP-derived TEQ, in humans [27-30], and has therefore been considered not only a biomarker of PCBs, but of overall POPs. Moreover, the major DDT metabolite 1,1-dichloro-2,2-bis (p-chlorophenyl)-ethylene (p,p'-DDE) which has anti-androgenic effect and is found in high levels in serum, has also been used. In the studies, both biomarkers were used in their lipid adjusted form, i.e., the wet weight concentrations were standardized using the sum of serum concentrations of triglycerides, cholesterol and phospholipids [27].

Since many of the studies on the Swedish Fishermen's Families Cohort have been of a retrospective nature, a drawback is that the biomarkers describe current, rather than past, exposure. Therefore, a model to estimate past

exposure was derived [31]. This model assumed a constant consumption of fatty fish, and a constant background exposure. However, the model took into consideration the decreasing levels of POPs in fatty fish, the reduction of body burden due to lactation, and the half-life of the compound in humans. The model was validated for CB-153 in women by comparing current concentrations of CB-153 to concentrations obtained from bio-banked samples from a rubella screening performed years earlier. The model has since then been validated for male CB-153, as well as for both male and female p,p'-DDE [32]. For CB-153, the correlations between biobanked (i.e. true) and estimated concentrations were 0.96 for men and 0.97 for women. When trichotomizing the female CB-153 concentrations, the kappa agreement between biobanked and estimated concentrations was 0.47 [33]. For p,p'-DDE the correlation between biobanked and estimated concentrations was 0.78 for both men and women [32].

The median plasma concentration of CB-153 among the fishermen's wives from the Swedish east coast was 160 ng/g lipid (range 16–780) and the median serum concentration among the sisters to the fishermen from the Swedish east coast was 115 ng/g lipid (range 27–560). With respect to the estimated past exposures, this varied from study to study depending on the time for which the concentrations were estimated, and which parameters were used in the

model. In those studies where estimated past exposure was related to any outcome, the estimated past concentrations was between 36% and 124% higher than the measured. The median serum concentrations among the fishermen were 193 ng/g lipid (range 40–1460) for CB-153 and 240 ng/g lipid (range 40–2250) for p,p'-DDE.

#### **Male reproduction parameters**

In the studies concerning male reproductive parameters, serum concentrations of CB-153 and p,p'-DDE were analyzed both as continuous measures (untransformed or log transformed) and categorized into five equally sized groups: For CB-153 the cut-points were 112, 167, 232, and 328 ng/g lipid, and for p,p'-DDE 136, 191, 273, and 471 ng/g lipid [16,19-21]. The correlation between CB-153 and p,p'-DDE for exposure was strong ( $r = 0.78$ ). Thus, the two biomarkers were never used simultaneously in any model.

A negative association, although not statistically significant when age was taken into account, between serum concentrations of CB-153 and the proportion of motile sperm was found [16]. Similar results were found for p,p'-DDE. No consistent associations were found between any of the two biomarkers and semen volume, sperm concentration, total sperm count or sperm morphology [16,34].

Sperm Chromatin Structure Assay (SCSA) was used to detect DNA fragmentation (small breaks in the sperm chromosome) [35,36]. The DNA fragmentation index (%DFI) was used to express the percentage of sperms showing DNA fragmentation [20]. Men in the lowest CB-153 quintile had lower %DFI than the men with higher CB-153 concentrations [20]. The pattern was similar for p,p'-DDE. With respect to continuous measures of CB-153 and p,p'-DDE there were no consistent association with %DFI.

Two color fluorescence in situ hybridization (FISH) was used to assess sperm chromosome Y:X ratio in an ejaculate [21]. Both CB-153 and p,p'-DDE were found to associate with an increased Y chromosome fraction [21,37]. When comparing the lowest exposed quintile to the highest, the effect was more marked for p,p'-DDE than for CB-153 [21]. Fructose, zinc, prostate specific antigen (PSA), and neutral  $\alpha$ -glucosidase (NAG) were analyzed as markers for the function of the epididymis, prostate and seminal vesicles [19]. Among these outcomes, the only significant association was found between CB-153 and total PSA [19].

When age was taken into account, no significant associations were observed between the POP biomarkers and serum hormone levels (FSH, LH, estradiol, testosterone,

SHBG, inhibin B, and the ratio testosterone/SHBG, respectively) [16].

#### **Female reproduction parameters**

East coast fishermen's wives and fishermen's sisters, who had grown up in a fishing village and/or fisherman's family tended to be somewhat older at menarche than referents from the same coastal area [38]. However, no differences were found between these women and the women from the west coast.

When adjusting for smoking, east coast fishermen's wives and sisters had somewhat shorter menstrual cycles than the west coast women. Moreover, in the east coast cohorts, women with a high consumption of fatty fish had shorter menstrual cycles than those with no or low fish consumption. However, there were no difference between women who had and had not grown up in a fishing village and/or fisherman's family, neither was there any effect of serum/plasma concentrations of CB-153.

There were no increased risk for miscarriage and stillbirths among the east coast fishermen's wives and fishermen's sisters, as compared to the corresponding cohorts from the west coast [25,39,40].

#### **Couple fertility**

A decreased fertility (i.e. prolonged TTP) was found for the east coast fishermen's wives as compared to the west coast fishermen's wives (Cox regression; Success Rate Ratio [SuRR] 0.86; 95% CI: 0.75, 0.99) [24]. The effect was most apparent among women who smoked more than 10 cigarettes per day, i.e. comparing east and west coast smokers (SuRR 0.68; 95% CI: 0.51, 0.91). Among the fishermen's sisters, no effect was found for cohort affiliation (logistic regression; Fecundability Odds Ratio [FOR] 0.99; 95% CI: 0.87, 1.14) [25]. The effect among smokers was not apparent in the group of fishermen's sisters. Among the east coast women, growing up in fishing village and/or fisherman's family did not suggest any negative effect on TTP, neither did a high consumption of fatty fish [24,25,39] or serum/plasma concentrations of CB-153 [26,33,39].

#### **Effects on the fetus**

Among the east coast fishermen's wives, the median birth weight was found to be 3530 g, which was lower than the corresponding figure for the west coast fishermen's wives (3610 g;  $p < 0.001$ ) [14]. The absolute birth weight difference was slightly lower among the fishermen's sisters (3500 g for the east coast women and 3560 g for the west coast women), although the statistical testing still found it significant ( $p < 0.001$ ) [17].

In a separate case-control analysis, a high total current intake of fish from the Baltic Sea indicated an increased risk of having an infant with low birth weight (OR = 1.9; 95% CI: 0.9, 3.9), although no dose-response relation was found [41]. Furthermore, when analyzing estimated intake of fish during the time period of interest (i.e. when the infant was born), the effects were diminished. However, childhood exposure (measured as growing up in a fishing village) carried an increased risk of giving birth to a low birth weight infant (OR = 2.1; 95% CI: 1.0, 4.3). The case mothers had higher median concentrations of CB-153 than the control mothers (190 ng/g lipid versus 160 ng/g lipid) [31].

There were no differences in birth length between the cohorts [14,17]. Among the fishermen's wives as well as the fishermen's sisters, the distribution of head circumference of the east coast infants was shifted downwards compared to the west coast infants ( $p < 0.001$  for both groups of women). An increased risk of being small for gestational age (SGA) [42] was found among infants born to the east coast fishermen's wives and sisters (OR = 1.5; 95% CI: 1.2, 2.0) [17]. Overall, there were no clear deviating patterns regarding gender ratio [14,17]. No specific malformation was overrepresented in the east coast cohorts [17,43].

### Discussion

The Swedish Fishermen's Families Cohorts consist of a large number of men and women of varying ages. The east coast cohorts comprises individuals who have been exposed to POPs at different times in their lives; those who are born and grown up in a fisherman's family and/or a fishing village are likely to have been exposed to POPs in utero and during their childhood, whereas others may have had high exposure only during their adult life. These assumptions are supported by that the men and women in the east coast cohorts have been found to have higher average levels of POPs than men and women from the general Swedish population [27,44,45], and those who have grown up in a fishing village and/or fishermen's family have been found to have especially high POP levels [45]. Thus, a major strength with the Swedish Fishermen's Families studies is the use of a large and relatively highly exposed study population.

The men and women in the east coast Swedish Fishermen's Families Cohorts have been found to eat more fish than individuals from the general Swedish population [14,17,22]. Since fish have constituents, e.g. long-chain n-3 fatty acids, minerals and vitamins, which may have positive effects on reproductive health [46-48], the general population may not be a suitable reference group in studies of health effects of dietary exposure to POPs. However, the waters off the west coast of Sweden are considerably

less contaminated than the Baltic Sea [15]. The west coast Swedish Fishermen's Families Cohorts have been found to eat fatty fish at a rate comparable to that of the east coast cohorts [14,22], but to have considerably lower levels of POPs [12]. Geographically, Sweden is a narrow country, only approximately 500 km at the widest location. Of the 9.1 million population, only 17% have non-Swedish origin [49]. The largest city, Stockholm, is inhabited by slightly more than 1 million individuals, making it a rather small city in an international perspective. All these factors taken together suggest that the Swedish population is a comparably homogenous one. With respect to the Swedish Fishermen's Families, the east and west coast cohorts, although residing at different coast lines, are geographically close. Moreover, although the east coast fishermen's wives tend to smoke more than those in the west coast cohorts [22,40], the west coast cohorts have been found to be similar to the east coast cohorts with respect to other socioeconomic factors, such as alcohol consumption [14,17,22] and educational level [17,40]. Hence, another major strength with the Swedish Fishermen's Families studies is the use of large and socio-economically similar cohorts as referent groups.

In several of the studies performed on the Swedish Fishermen's Families Cohorts, biomarkers for exposure to POPs have been used. Most frequently, CB-153 has been considered as a biomarker for a person's total body burden of POP. The rationale for this is that this PCB congener has been found to correlate strongly with the total PCB concentration in plasma [27], serum [28], whole venous blood [50], and cord blood [50], as well as with the TEQ from total PCB [27,30] and total POP-derived TEQ [6]. Other proxy measures of exposure used in the Swedish Fishermen's Families Cohorts, i.e. cohort affiliation, fish consumption and grown up in a fisherman's family and/or fishing village, have been found to be strongly associated with the concentrations of CB-153 [12,27,45]. Thus, a further strength with the Swedish Fishermen's Families Cohorts studies is the use of biomarkers of exposure. However, the studies are in general retrospective or cross-sectional and the proper exposure window might not always be reflected in an optimal way. For instance, in the study which investigated the association between POP exposure and low birth weight, we had to estimate the concentrations of CB-153 during year of childbirth based on back-calculation models [31]. This might lead to a negative impact on precision, as well as accuracy, of the estimated CB-153 concentrations. As stated by Grandjean et al [51], it is probably not only the dose that makes the poison, of equally importance might be the timing of exposure. Accordingly, future studies should force to collect exposure information that in a better way reflect the most relevant exposure window.

When they enter the human body, POPs are stored in the adipose tissue. Thus, to obtain a valid measure of the body burden of POP, a biopsy should be performed and fatty tissue should be analyzed. This is, however, a procedure which is more costly and painful than collecting a blood sample. Nevertheless, it may be assumed that the concentration of POP in serum lipids is in equilibrium with that in adipose tissue. Kahn et al [52] found a very high correlation ( $r = 0.89$ ) between 2,3,7,8 TCDD concentration in fat biopsy and the lipid adjusted serum concentration. Thus, the body burden of POP may be estimated by calculating the lipid standardized serum concentrations. However, Schisterman et al [53] have recently shown that this method is highly prone to bias, causing an underestimation of the true effect. Nevertheless, the correlations between wet weight and lipid adjusted concentrations of POPs in the Swedish Fishermen's Families Cohorts are high [27,45]. Moreover, the results obtained when using the biomarkers are in agreement with those obtained using the proxy measures of exposure. Thus, a possible bias introduced by the lipid standardization should pose, at most, a minor problem.

A variety of outcomes have been investigated in the Swedish Fishermen's Families Cohort. The ones chosen have been such that they are believed to cover all aspects of human reproduction, i.e. parameters concerning the man, the woman, the couple and the fetus. By using this multitude of outcomes, we have been able to establish where POPs may do most harm to human reproduction, i.e. to the fetus. However, the retrospective design may pose a problem for some outcomes, where recall was long and a possible misclassification may have been introduced. The most apparent example of this is menarche. For other outcomes, mainly those related to the fetus, data has been collected from registries and the quality should therefore be higher. Also, the studies concerning male reproductive factors were not affected by any recall bias.

The most consistent results from the Swedish Fishermen's Families Cohorts are those on infant parameters, and especially birth weight and fetal growth. Irrespective of which measure of exposure was used, children who are exposed to POPs in utero have an increased risk of weighing less at birth, also after having taken the gestational length into account (SGA). Similar findings have been observed among fish eaters from the Lake Michigan area in the US who had PCB levels comparable to the ones among the Swedish fishermen's wives [54]. Also in the general population in the Netherlands with relatively low POP exposures, in utero levels of PCB were negatively associated with birth weight [55]. There are, however, also epidemiological examples showing no, or even positive, associations between POP exposure and birth weight [56-58]. The overall picture regarding the hypothesized asso-

ciation between POP exposure and birth weight is, accordingly, still far from conclusive.

Environmental chemicals acting as sex hormone agonists or antagonists could have an adverse effect on male sex hormones which in turn might decrease the male reproductive function [4,59]. In our study sperm motility decreased with increasing serum levels of POP. This finding is consistent with four other cross-sectional studies carried out during recent years [60-63]. Sperm DNA integrity is essential for the accurate transmission of genetic information and sperm chromatin abnormalities or DNA damage may result in impaired male fertility [64]. Among Swedish fishermen we found that increasing POP exposure increased the level of sperm DNA strand breaks. However, in a study comprising Inuits, with higher serum levels of POP, the levels of sperm DNA strandbreaks were low. This inconsistent result might be explained by differences in the genetic background and lifestyle habits, which still needs to be elucidated. The mechanism behind changes in Y:X ratio is unknown. However one hypothesis could be a loss of X chromosome due to an effect on formation of micronuclei during the meiosis, by *e.g.* POPs. We found effects of POP biomarkers on the proportion of Y- and X-bearing sperm. However, such effects were not seen in three other study populations [65]. The inconsistent results might be explained by differences in POP exposure profile and dose due to variation in lifestyle and diet. Regarding sperm count and sperm morphology none of these outcomes were related to POP exposure.

The results concerning female reproductive parameters are ambiguous, and in some cases they suggest a protective, rather than hazardous, effect of fish consumption. Taking other studies on the topic into account, no obvious risk pattern emerges for most of the outcomes. Although results from a cross-sectional study on Akwesasne Mohawk girls suggested that high concentrations of blood PCB lead to a higher probability of early menarche [66], retrospective studies performed by Vasiliu et al [67] and in the Swedish Fishermen's Families Cohort studies found no such effect. Moreover, whereas exposure to POPs was associated with a shortened menstrual cycle length in the retrospective studies performed in the Swedish Fishermen's Families Cohorts and the New York State Angler Cohort [68], another retrospective study found the opposite relation between exposure and outcome [69], whereas a prospective study using daily urine sampling found no relation at all [70]. With respect to miscarriages and stillbirths, the results available are more homogenous than those for the other outcomes. In the Swedish Fishermen's Families Cohorts, none of the analyses produced a statistically significant hazard associated with POP exposure. However, in some of the subgroup analyses, POP exposure was suggested to have a protective effect. This is in

agreement with other studies, were no effect [71-73], or a protective effect [74], has been found.

East coast fishermen's wives were found to have longer TTPs than west coast fishermen's wives. However, these results were not replicated among the fishermen's sisters, neither did any of the exposure measures within the east coast cohorts suggest any effect of POP exposure. In the New York State Angler Cohort, several studies have been performed regarding parental POP exposure and fertility. The majority of these results show no effect on TTP, delayed conception, or infertility of maternal [75,76] or paternal [77,78] exposure, although one study did suggest reduced fecundability among women with high consumption of fatty fish from the Great Lakes [78]. In the INUENDO study, results suggested that high exposure to POPs were associated with reduced fertility only among Inuits from Greenland, and not among couples, i.e. maternal and paternal exposure, from Poland, Ukraine or Sweden [79].

When evaluating the results from the Swedish Fishermen's Families Cohorts studies, it is important to keep in mind that the main exposure source for POP, namely fatty fish from the Baltic Sea, is also an exposure route for other constituents of fish. These other constituents include both other pollutants, such as methylmercury, and natural components, such as marine n-3 fatty acids, minerals and vitamins. In the Swedish Fishermen's Families Cohorts, the concentration of CB-153 has been found to be correlated ( $r_s = 0.51$ ) to that of methylmercury [80]. However, where associations were found between on one hand CB-153 and p,p'-DDE and on the other hand several male reproductive parameters, no consistent relationships were found between methylmercury and the same male parameters.

With respect to the natural components of fatty fish, marine n-3 fatty acids have been found to be associated with prolonged gestation in randomized trials [81-85]. Moreover, positive associations between seafood consumption and reproductive outcomes, such as birth weight and gestational length, have been seen in observational studies [47,48,86-90]. Thus, there is a possibility that the effect of the potentially hazardous constituents in Baltic Sea fatty fish could be outweighed by their beneficial counterparts, which would explain the positive results found for some outcomes.

## Conclusion

The major strengths of the studies were that the study populations included socio-economic similar cohorts, which enabled good opportunities to control for confounding, and the cohorts included the necessary exposure contrasts, which made it possible to investigate the hypothe-

sized association between POP exposure and reproductive health.

The main finding from the studies performed on the Swedish Fishermen's Families Cohorts is that although there may be minor effects on both the male and female reproduction system, the major effects are those concerning the growth of the fetus. This observation was strengthened by the fact that the negative effect on birth weight was observed for all exposure measures used. In addition, corresponding effects have been observed in other populations with a similar exposure source, i.e. intake of contaminated fish.

Regarding male and female reproductive data, the Swedish data together with other epidemiological studies, provide no conclusive results. Although some associations may have been affected by beneficial effects of essential nutrients in seafood, the overall findings are meaningful in the context of reproductive toxicity and support the usefulness of the epidemiological design.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

AA initiated the project. All authors participated in drafting the manuscript, and have approved of the final version.

## Acknowledgements

The studies on the Swedish Fishermen's Families Cohort were all carried out in collaboration with Professor Lars Hagmar, who passed away in June 2006.

## References

1. Jensen S: **The PCB story.** *Ambio* 1972, **1**:123-131.
2. Colborn T, vom Saal FS, Soto AM: **Developmental effects of endocrine-disrupting chemicals in wildlife and humans.** *Environ Health Perspect* 1993, **101**(5):378-384.
3. Kelce WR, Stone CR, Laws SC, Gray LE, Kempainen JA, Wilson EM: **Persistent DDT metabolite p,p'-DDE is a potent androgen receptor antagonist.** *Nature* 1995, **375**(6532):581-585.
4. Sharpe RM, Skakkebaek NE: **Are oestrogens involved in falling sperm counts and disorders of the male reproductive tract?** *Lancet* 1993, **341**(8857):1392-1395.
5. AMAP: **Arctic pollution issues.** Oslo: Arctic monitoring and assessment programme (AMAP); 1998.
6. Brouwer A, Ahlborg UG, Berg M Van den, Birnbaum LS, Boersma ER, Bosveld B, Denison MS, Gray LE, Hagmar L, Holene E, Huisman M, Jacobson SW, Jacobson JL, Koopman-Esseboom C, Koppe JG, Kulig BM, Morse DC, Muckle G, Peterson RE, Sauer PJJ, Seegal RF, Smits-Van Prooije AE, Touwen BCL, Weisglas-Kuperus N, Winneke G: **Functional aspects of developmental toxicity of polyhalogenated aromatic hydrocarbons in experimental animals and human infants.** *Eur J Pharmacol* 1995, **293**(1):1-40.
7. Fångström B, Hovander L, Bignert A, Athanassiadis I, Linderholm L, Grandjean P, Weihe P, Bergman Å: **Concentrations of polybrominated diphenyl ethers, polychlorinated biphenyls, and polychlorobiphenyls in serum from pregnant Faroese women and their children 7 years later.** *Environ Sci Technol* 2005, **39**(24):9457-9463.

8. Fångström B, Strid A, Grandjean P, Weihe P, Bergman Å: **A retrospective study of PBDEs and PCBs in human milk from the Faroe Islands.** *Environ Health* 2005, **4**:12.
9. Jönsson B, Rylander L, Lindh C, Rignell-Hydbom A, Giwercman A, Toft G, Pedersen HS, Ludwicki JK, Goralczyk K, Zvezday V, Spanò M, Bizzaro D, Bonefeld-Jørgensen EC, Manicardi GC, Bonde JP, Hagmar L: **Inter-population variations in concentrations, determinants of and correlations between 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) and 1,1-dichloro-2,2-bis (p-chlorophenyl)-ethylene (p,p'-DDE): a cross-sectional study of 3161 men and women from Inuit and European populations.** *Environ Health* 2005, **4**:27.
10. Sjödin A, Jones RS, Focant JF, Lapeza C, Wang RY, McGahee EE 3rd, Zhang Y, Turner WE, Slazyk B, Needham LL, Patterson DG Jr: **Retrospective time-trend study of polybrominated diphenyl ether and polybrominated and polychlorinated biphenyl levels in human serum from the United States.** *Environ Health Perspect* 2004, **112**(6):654-658.
11. Fangstrom B, Athanasiadou M, Grandjean P, Weihe P, Bergman A: **Hydroxylated PCB metabolites and PCBs in serum from pregnant Faroese women.** *Environ Health Perspect* 2002, **110**(9):895-899.
12. Svensson BG, Nilsson A, Hansson M, Rappe C, Akesson B, Skerfving S: **Exposure to dioxins and dibenzofurans through the consumption of fish.** *N Engl J Med* 1991, **324**:8-12.
13. Svensson BG, Mikoczy Z, Stromberg U, Hagmar L: **Mortality and cancer incidence among Swedish fishermen with a high dietary intake of persistent organochlorine compounds.** *Scand J Work Environ Health* 1995, **21**:106-115.
14. Rylander L, Stromberg U, Hagmar L: **Decreased birthweight among infants born to women with a high dietary intake of fish contaminated with persistent organochlorine compounds.** *Scand J Work Environ Health* 1995, **21**:368-375.
15. Bergqvist P, Bergek S, Hallbäck H, Rappe C, Slorach S: **Dioxins in cod and herring from the seas around Sweden.** *Chemosphere* 1989, **19**:513-516.
16. Rignell-Hydbom A, Rylander L, Giwercman A, Jönsson BA, Nilsson-Ehle P, Hagmar L: **Exposure to CB-153 and p,p'-DDE and male reproductive function.** *Hum Reprod* 2004, **19**:2066-2075.
17. Rylander L, Stromberg U, Hagmar L: **Lowered birth weight among infants born to women with a high intake of fish contaminated with persistent organochlorine compounds.** *Chemosphere* 2000, **40**:1255-1262.
18. Hagmar L, Linden K, Nilsson A, Norrving B, Akesson B, Schutz A, Moller T: **Cancer incidence and mortality among Swedish Baltic Sea fishermen.** *Scand J Work Environ Health* 1992, **18**(4):217-224.
19. Rignell-Hydbom A, Rylander L, Elzanaty S, Giwercman A, Lindh CH, Hagmar L: **Exposure to persistent organochlorine pollutants and seminal levels of markers of epididymal and accessory sex gland functions in Swedish men.** *Hum Reprod* 2005, **20**(7):1910-1914.
20. Rignell-Hydbom A, Rylander L, Giwercman A, Jonsson BA, Lindh C, Eleuteri P, Rrescia M, Leter G, Cordelli E, Spano M, Hagmar L: **Exposure to PCBs and p,p'-DDE and human sperm chromatin integrity.** *Environ Health Perspect* 2005, **113**:175-179.
21. Tiido T, Rignell-Hydbom A, Jonsson B, Giwercman YL, Rylander L, Hagmar L, Giwercman A: **Exposure to persistent organochlorine pollutants associates with human sperm Y:X chromosome ratio.** *Hum Reprod* 2005, **20**:1903-1909.
22. Rylander L, Hagmar L: **Mortality and cancer incidence among women with a high consumption of fatty fish contaminated with persistent organochlorine compounds.** *Scand J Work Environ Health* 1995, **21**(1):419-426.
23. Rylander L, Stromberg U, Hagmar L: **Agreement between reported fish consumption obtained by two interviews and its impact on the results in a reproduction study.** *Eur J Epidemiol* 1998, **14**:93-97.
24. Axmon A, Rylander L, Strömberg U, Hagmar L: **Time to pregnancy and infertility among women with a high intake of fish contaminated with persistent organochlorine compounds.** *Scand J Work Environ Health* 2000, **26**:199-206.
25. Axmon A, Rylander L, Strömberg U, Hagmar L: **Female fertility in relation to the consumption of fish contaminated with persistent organochlorine compounds.** *Scand J Work Environ Health* 2002, **28**(2):124-132.
26. Axmon A, Rylander L, Strömberg U, Jönsson B, Nilsson-Ehle P, Hagmar L: **Polychlorinated biphenyls in serum and time to pregnancy.** *Environ Res* 2004, **96**(2):186-195.
27. Grimvall E, Rylander L, Nilsson-Ehle P, Nilsson U, Strömberg U, Hagmar L, Östman C: **Monitoring of polychlorinated biphenyls in human blood plasma: methodological developments and influence of age, lactation, and fish consumption.** *Arch Environ Contam Toxicol* 1997, **32**(3):329-336.
28. Glynn AW, Wolok A, Aune M, Atuma S, Zettermark S, Maehle-Schmid M, Darnerud PO, Becker W, Vessby B, Adami HO: **Serum concentrations of organochlorines in men: a search for markers of exposure.** *Sci Total Environ* 2000, **263**(1-3):197-208.
29. Hagmar L, Becher G, Heikkilä A, Frankman O, Dyremark E, Schutz A, Ahlborg UG, Dybing E: **Consumption of fatty fish from the Baltic Sea and PCB in whole venous blood, plasma and cord blood from delivering women in the Åland/Turku archipelago.** *J Toxicol Environ Health A* 1998, **53**(8):581-591.
30. Glynn AW, Atuma S, Aune M, Darnerud PO, Cnattingius S: **Polychlorinated biphenyl congeners as markers of toxic equivalents of polychlorinated biphenyls, dibenzo-p-dioxins and dibenzofurans in breast milk.** *Environ Res* 2001, **86**(3):217-228.
31. Rylander L, Stromberg U, Dyremark E, Ostman C, Nilsson-Ehle P, Hagmar L: **Polychlorinated biphenyls in blood plasma among Swedish female fish consumers in relation to low birth weight.** *Am J Epidemiol* 1998, **147**(5):493-502.
32. Axmon A, Rignell-Hydbom A: **Estimations of past male and female serum concentrations of biomarkers of persistent organochlorine pollutants and their impact on fecundability estimates.** *Environ Res* 2006, **101**(3):87-94.
33. Axmon A, Rylander L, Strömberg U, Dyremark E, Hagmar L: **Polychlorinated biphenyls in blood plasma among Swedish female fish consumers in relation to time to pregnancy.** *J Toxicol Environ Health A* 2001, **64**(6):485-498.
34. Toft G, Rignell-Hydbom A, Tyrkiel E, Shvets M, Giwercman A, Lindh CH, Pedersen HS, Ludwicki JK, Lesovoy V, Hagmar L, Spano M, Manicardi GC, Bonefeld-Jørgensen EC, Thulstrup AM, Bonde JP: **Semen quality and exposure to persistent organochlorine pollutants.** *Epidemiology* 2006, **17**(4):450-458.
35. Evenson DP, Larson KL, Jost LK: **Sperm chromatin structure assay: its clinical use for detecting sperm DNA fragmentation in male infertility and comparisons with other techniques.** *J Androl* 2002, **23**(1):25-43.
36. Spano M, Bonde JP, Hjollund HI, Kolstad HA, Cordelli E, Leter G: **Sperm chromatin damage impairs human fertility. The Danish First Pregnancy Planner Study Team.** *Fertil Steril* 2000, **73**(1):43-50.
37. Vittinghoff E, Glidden D, Shiboski S, McCulloch C: **Regression Methods in Biostatistics - Linear, Logistic, Survival, and Repeated Measures Models.** Springer 2005.
38. Axmon A: **Menarche in women with high exposure to persistent organochlorine pollutants in utero and during childhood.** *Environ Res* 2006, **102**(1):77-82.
39. Axmon A: **Fertility and female dietary exposure to persistent organochlorine compounds.** In *Doctoral thesis* Lund: Department of Occupational and Environmental Medicine, Institute of Laboratory Medicine, Lund University; 2003.
40. Axmon A, Rylander L, Strömberg U, Hagmar L: **Miscarriages and stillbirths in women with a high intake of fish contaminated with persistent organochlorine compounds.** *Int Arch Occup Environ Health* 2000, **73**:204-208.
41. Rylander L, Stromberg U, Hagmar L: **Dietary intake of fish contaminated with persistent organochlorine compounds in relation to low birthweight.** *Scand J Work Environ Health* 1996, **22**(4):260-266.
42. Källén B: **A birth weight for gestational age standard based on data in the Swedish Medical Birth Registry, 1985-1989.** *Eur J Epidemiol* 1995, **11**(5):601-606.
43. Rylander L, Hagmar L: **No evidence for congenital malformations or prenatal death in infants born to women with a high dietary intake of fish contaminated with persistent organochlorines.** *Int Arch Occup Environ Health* 1999, **72**:121-124.
44. Svensson BG, Nilsson A, Jonsson E, Schütz A, Åkesson B, Hagmar L: **Fish consumption and exposure to persistent organochlorine compounds, mercury, selenium and methylamines among Swedish fishermen.** *Scand J Work Environ Health* 1995, **21**(2):96-105.

45. Rylander L, Dyremark E, Stromberg U, Ostman C, Hagmar L: **The impact of age, lactation and dietary habits on PCB in plasma in Swedish women.** *Sci Total Environ* 1997, **207(1)**:55-61.
46. Olsen SF: **Consumption of marine n-3 fatty acids during pregnancy as a possible determinant of birth weight. A review of the current epidemiologic evidence.** *Epidemiol Rev* 1993, **15**:399-413.
47. Olsen SF, Grandjean P, Weihe P, Videro T: **Frequency of seafood intake in pregnancy as a determinant of birth weight: evidence for a dose dependent relationship.** *J Epidemiol Community Health* 1993, **47**:436-440.
48. Odent MR, McMillan L, Kimmel T: **Prenatal care and sea fish.** *Eur J Obstet Gynecol Reprod Biol* 1996, **68**:49-51.
49. **Statistics Sweden** [<http://www.scb.se>]
50. Hagmar L, Becher G, Heikkilä A, Frankman O, Dyremark E, Schütz A, Ahlborg UG, Dybing E: **Consumption of fatty fish from the Baltic Sea and PCB in whole venous blood, plasma and cord blood from delivering women in the Åland/Turku archipelago.** *J Toxicol Environ Health A* 1998, **53(8)**:581-591.
51. Grandjean P, Bellinger D, Bergman A, Cordier S, Davey-Smith G, Eskenazi B, Gee D, Gray K, Hanson M, Hazel P van den, Heindel JJ, Heinzow B, Hertz-Picciotto I, Hu H, Huang TT, Jensen TK, Landrigan PJ, McMillan IC, Murata K, Ritz B, Schoeters G, Skakkebaek NE, Skerfving S, Weihe P: **The faroes statement: human health effects of developmental exposure to chemicals in our environment.** *Basic Clin Pharmacol Toxicol* 2008, **102(2)**:73-75.
52. Kahn PC, Gochfeld M, Nygren M, Hansson M, Rappe C, Velez H, Ghent-Guenther T, Wilson WP: **Dioxins and dibenzofurans in blood and adipose tissue of Agent Orange-exposed Vietnam veterans and matched controls.** *JAMA* 1988, **259(11)**:1661-1667.
53. Schisterman EF, Whitcomb BW, Louis GM, Louis TA: **Lipid adjustment in the analysis of environmental contaminants and human health risks.** *Environ Health Perspect* 2005, **113(7)**:853-857.
54. Fein GG, Jacobson JL, Jacobson SW, Schwartz PM, Dowler JK: **Prenatal exposure to polychlorinated biphenyls: effects on birth size and gestational age.** *J Pediatr* 1984, **105(2)**:315-320.
55. Patandin S, Koopman-Esseboom C, de Ridder MA, Weisglas-Kuperus N, Sauer PJ: **Effects of environmental exposure to polychlorinated biphenyls and dioxins on birth size and growth in Dutch children.** *Pediatr Res* 1998, **44(4)**:538-545.
56. Vartiainen T, Jaakkola JJ, Saarikoski S, Tuomisto J: **Birth weight and sex of children and the correlation to the body burden of PCDDs/PCDFs and PCBs of the mother.** *Environ Health Perspect* 1998, **106(2)**:61-66.
57. Rogan WJ, Gladen BC, McKinney JD, Carreras N, Hardy P, Thullen J, Tingelstad J, Tully M: **Polychlorinated biphenyls (PCBs) and dichlorodiphenyl dichloroethene (DDE) in human milk: effects on growth, morbidity, and duration of lactation.** *Am J Public Health* 1987, **77(10)**:1294-1297.
58. Dar E, Kanarek MS, Anderson HA, Sonzogni WC: **Fish consumption and reproductive outcomes in Green Bay, Wisconsin.** *Environ Res* 1992, **59(1)**:189-201.
59. Toppari J, Larsen JC, Christiansen P, Giwercman A, Grandjean P, Guillelte LJ Jr, Jegou B, Jensen TK, Jouannet P, Keiding N, Leffers H, McLachlan JA, Meyer O, Muller J, Rajpert-De Meyts E, Scheike T, Sharpe R, Sumpster J, Skakkebaek NE: **Male reproductive health and environmental xenoestrogens.** *Environ Health Perspect* 1996, **104(Suppl 4)**:741-803.
60. Guo YL, Hsu PC, Hsu CC, Lambert GH: **Semen quality after prenatal exposure to polychlorinated biphenyls and dibenzofurans.** *Lancet* 2000, **356(9237)**:1240-1241.
61. Dallinga JW, Moonen EJ, Dumoulin JC, Evers JL, Geraedts JP, Kleinjans JC: **Decreased human semen quality and organochlorine compounds in blood.** *Hum Reprod* 2002, **17**:1973-1979.
62. Richthoff J, Rylander L, Jönsson BA, Åkesson H, Hagmar L, Nilsson-Ehle P, Stridsberg M, Giwercman A: **Serum levels of 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) in relation to markers of reproductive function in young males from the general Swedish population.** *Environ Health Perspect* 2003, **111(4)**:409-413.
63. Rozati R, Reddy PP, Reddanna P, Mujtaba R: **Role of environmental estrogens in the deterioration of male factor fertility.** *Fertil Steril* 2002, **78(6)**:1187-1194.
64. Agarwal A, Said TM: **Role of sperm chromatin abnormalities and DNA damage in male infertility.** *Hum Reprod Update* 2003, **9(4)**:331-345.
65. Tiido T, Rignell-Hydbom A, Jonsson BA, Giwercman YL, Pedersen HS, Wojtyniak B, Ludwicki JK, Lesovoy V, Zvyezday V, Spano M, Manicardi GC, Bizzaro D, Bonefeld-Jorgensen EC, Toft G, Bonde JP, Rylander L, Hagmar L, Giwercman A: **Impact of PCB and p,p'-DDE contaminants on human sperm Y:X chromosome ratio: studies in three European populations and the Inuit population in Greenland.** *Environ Health Perspect* 2006, **114**:718-724.
66. Denham M, Schell LM, Deane G, Gallo MV, Ravenscroft J, DeCaprio AP: **Relationship of lead, mercury, mirex, dichlorodiphenyl-dichloroethylene, hexachlorobenzene, and polychlorinated biphenyls to timing of menarche among Akwesasne Mohawk girls.** *Pediatrics* 2005, **115(2)**:e127-134.
67. Vasiliu O, Muttineni J, Karmaus W: **In utero exposure to organochlorines and age at menarche.** *Hum Reprod* 2004, **19(7)**:1506-1512.
68. Mendola P, Buck GM, Sever LE, Zielezny M, Vena JE: **Consumption of PCB-contaminated freshwater fish and shortened menstrual cycle length.** *Am J Epidemiol* 1997, **146(11)**:955-960.
69. Cooper GS, Klebanoff MA, Promislow J, Brock JW, Longnecker MP: **Polychlorinated biphenyls and menstrual cycle characteristics.** *Epidemiology* 2005, **16(2)**:191-200.
70. Windham GC, Lee D, Mitchell P, Anderson M, Petreas M, Lasley B: **Exposure to organochlorine compounds and effects on ovarian function.** *Epidemiology* 2005, **16**:182-190.
71. Khanjani N, Sim MR: **Maternal contamination with PCBs and reproductive outcomes in an Australian population.** *J Expo Sci Environ Epidemiol* 2006.
72. Khanjani N, Sim MR: **Maternal contamination with dichlorodiphenyltrichloroethane and reproductive outcomes in an Australian population.** *Environ Res* 2006, **101(3)**:373-379.
73. Sugiura-Ogasawara M, Ozaki Y, Sonta S, Makino T, Suzumori K: **PCBs, hexachlorobenzene and DDE are not associated with recurrent miscarriage.** *Am J Reprod Immunol* 2003, **50**:485-489.
74. Mendola P, Buck GM, Vena JE, Zielezny M, Sever LE: **Consumption of PCB-contaminated sport fish and risk of spontaneous fetal death.** *Environ Health Perspect* 1995, **103(5)**:498-502.
75. Buck GM, Sever LE, Mendola P, Zielezny M, Vena JE: **Consumption of contaminated sport fish from Lake Ontario and time-to-pregnancy. New York State Angler Cohort.** *Am J Epidemiol* 1997, **146(11)**:949-954.
76. McGuinness BM, Buck GM, Mendola P, Sever LE, Vena JE: **Infecundity and consumption of polychlorinated biphenyl-contaminated fish.** *Arch Environ Health* 2001, **56(3)**:250-253.
77. Buck GM, Mendola P, Vena JE, Sever LE, Kostyniak P, Greizerstein H, Olson J, Stephen FD: **Paternal Lake Ontario fish consumption and risk of conception delay. New York State Angler Cohort.** *Environ Res* 1999, **80(2Pt2)**:S13-S18.
78. Buck GM, Vena JE, Schisterman EF, Dmochowski J, Mendola P, Sever LE, Fitzgerald E, Kostyniak P, Greizerstein H, Olson J: **Parental consumption of contaminated sport fish from Lake Ontario and predicted fecundability.** *Epidemiology* 2000, **11(4)**:388-393.
79. Axmon A, Thulstrup AM, Rignell-Hydbom A, Pedersen HS, Zvyezday V, Ludwicki JK, Jonsson BA, Toft G, Bonde JP, Hagmar L: **Time to pregnancy as a function of male and female serum concentrations of 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) and 1,1-dichloro-2,2-bis (p-chlorophenyl)-ethylene (p,p'-DDE).** *Hum Reprod* 2006, **21**:657-665.
80. Rignell-Hydbom A, Axmon A, Lundh T, Jonsson BA, Tiido T, Spano M: **Dietary exposure to methyl mercury and PCB and the associations with semen parameters among Swedish fishermen.** *Environ Health* 2007, **6**:14.
81. Smuts CM, Huang M, Mundy D, Plasse T, Major S, Carlson SE: **A randomized trial of docosahexaenoic acid supplementation during the third trimester of pregnancy.** *Obstet Gynecol* 2003, **101(3)**:469-479.
82. Olsen SF, Secher NJ: **A possible preventive effect of low-dose fish oil on early delivery and pre-eclampsia: indications from a 50-year-old controlled trial.** *Br J Nutr* 1990, **64**:599-609.
83. Olsen SF, Secher NJ, Tabor A, Weber T, Walker JJ, Glud C: **Randomised clinical trials of fish oil supplementation in high risk pregnancies. Fish Oil Trials In Pregnancy (FOTIP) Team.** *Bjog* 2000, **107(3)**:382-395.

84. Olsen SF, Sorensen JD, Secher NJ, Hedegaard M, Henriksen TB, Hansen HS, Grant A: **Randomised controlled trial of effect of fish-oil supplementation on pregnancy duration.** *Lancet* 1992, **339(8800):1003-1007.**
85. Olsen SF, Osterdal ML, Salvig JD, Weber T, Tabor A, Secher NJ: **Duration of pregnancy in relation to fish oil supplementation and habitual fish intake: a randomised clinical trial with fish oil.** *Eur J Clin Nutr* 2007, **61(8):976-985.**
86. Olsen SF, Hansen HS, Sorensen TI, Jensen B, Secher NJ, Sommer S, Knudsen LB: **Intake of marine fat, rich in (n-3)-polyunsaturated fatty acids, may increase birthweight by prolonging gestation.** *Lancet* 1986, **2:367-369.**
87. Olsen SF, Olsen J, Frische G: **Does fish consumption during pregnancy increase fetal growth? A study of the size of the newborn, placental weight and gestational age in relation to fish consumption during pregnancy.** *Int J Epidemiol* 1990, **19(4):971-977.**
88. Olsen SF, Secher NJ: **Low consumption of seafood in early pregnancy as a risk factor for preterm delivery: prospective cohort study.** *Bmj* 2002, **324:447.**
89. Thorsdottir I, Birgisdottir BE, Halldorsdottir S, Geirsson RT: **Association of fish and fish liver oil intake in pregnancy with infant size at birth among women of normal weight before pregnancy in a fishing community.** *Am J Epidemiol* 2004, **160(5):460-465.**
90. Petridou E, Stoikidou M, Diamantopoulou M, Mera E, Dessypris N, Trichopoulos D: **Diet during pregnancy in relation to birthweight in healthy singletons.** *Child Care Health Dev* 1998, **24(3):229-242.**
91. Axmon A, Rylander L, Strömberg U, Hagmar L: **Altered menstrual cycles in women with a high dietary intake of persistent organochlorine compounds.** *Chemosphere* 2004, **56:813-819.**

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

