

A Multi-city Time-series Analysis of Air Pollution and Emergency Department Visits for
Cardiac and Respiratory Conditions

David M. Stieb (corresponding author), Population Studies Division, Healthy
Environments and Consumer Safety Branch, Health Canada, and Department of
Epidemiology and Community Medicine, University of Ottawa, Postal locator 4903C,
269 Laurier Ave. W., Ottawa, ON, Canada, K1A 0K9, 613 957 3132 (voice); 613 954
7612 (fax); dave_stieb@hc-sc.gc.ca

Mieczyslaw Szyszkowicz, Population Studies Division, Healthy Environments and
Consumer Safety Branch, Health Canada, Postal locator 4903C, 269 Laurier Ave. W.,
Ottawa, ON, Canada, K1A 0K9, mietek_szyszkowicz@hc-sc.gc.ca

Brian H. Rowe, Department of Emergency Medicine, University of Alberta, Room
1G1.43 WMC, University of Alberta Hospital, 8440-112th Street, Edmonton, Alberta,
CANADA T6G 2B7, brian.rowe@ualberta.ca

Judith A. Leech, Department of Medicine, University of Ottawa and Population Studies
Division, Healthy Environments and Consumer Safety Branch, Health Canada, Ottawa
Hospital, Civic Campus, 1053 Carling Avenue, Ottawa, Ontario, Canada, K1Y 4E9,
jleech@ottawahospital.on.ca

ABSTRACT

BACKGROUND: Relatively few studies have been conducted of the association between air pollution and emergency department (ED) visits, and most of these have been based on a small number of visits, for a limited number of health conditions and pollutants, and only daily measures of exposure and response.

METHODS: A time-series analysis was conducted on nearly 400,000 ED visits to 14 hospitals in seven Canadian cities. Associations were examined between carbon monoxide (CO), nitrogen dioxide (NO₂), ozone (O₃), sulfur dioxide (SO₂), and particulate matter of median aerodynamic diameter less than 10 and 2.5 microns (PM₁₀ and PM_{2.5}), and visits for angina/myocardial infarction, heart failure, dysrhythmia/conduction disturbance, asthma, chronic obstructive pulmonary disease (COPD), and respiratory infections. Daily and 3-hourly visit counts were modeled as quasi-Poisson and analyses controlled for effects of temporal cycles, weather, day of week and holidays.

RESULTS: 24-hour average concentrations of CO and NO₂ lag 0 days exhibited the most consistent associations with cardiac conditions (2.1% (95% CI 0.0-4.2%) and 2.6% (95% CI 0.2-5.0%) increase in visits for myocardial infarction/angina per 0.7 ppm CO and 18.4 ppb NO₂ respectively; 3.8% (95% CI 0.7-6.9%) and 4.7% (95% CI 1.2-8.4%) increase in visits for heart failure). Ozone (lag 2 days) was most consistently associated with respiratory visits (3.2% (95% CI 0.3-6.2%), and 3.7% (95% CI -0.5-7.9%) increases in asthma and COPD visits respectively per 18.4 ppb). Associations tended to be of greater magnitude during the warm season (April to September). In particular, the associations of PM₁₀ and PM_{2.5} with asthma visits were respectively nearly three- and over fourfold larger (14.4% increase in visits, 95% CI, 0.2-30.7, per 20.6 µg/ m³ PM₁₀ and 7.6%

increase in visits, 95% CI, 5.1-10.1, per 8.2 $\mu\text{g}/\text{m}^3$ $\text{PM}_{2.5}$). No consistent associations were observed between three hour averages of pollutant concentrations and ED visits.

CONCLUSIONS: In this large multicenter analysis, daily average concentrations of CO and NO_2 exhibited the most consistent associations with ED visits for cardiac conditions, while ozone exhibited the most consistent associations with visits for respiratory conditions. PM_{10} and $\text{PM}_{2.5}$ were strongly associated with asthma visits during the warm season.

BACKGROUND

Numerous analyses have been conducted of the association between outdoor air pollution and mortality [1], as well as hospital admissions, owing to the wide availability of these data through national vital statistics and health care statistics systems. While these studies have been central to establishing the occurrence of adverse health effects of air pollution even at the relatively low levels of exposure now observed in most industrialized countries, these health outcomes are generally believed to occur in a relatively small segment of the population in comparatively poor health. The emergency department (ED) is often the point of entry for those ultimately admitted to hospital. For many cardiac and respiratory conditions, however, less than half (and as few as 10% in the case of asthma) of those seen in the emergency department are admitted to hospital [2]. Thus ED visits reflect impacts in a broader segment of the population.

Since these data are not routinely available, fewer studies have been conducted, and of these, most have been based on a single city and a relatively small number of visits, for a limited number of health conditions and pollutants. Multi-city studies are believed to generate more stable results which are less prone to biases that may affect small studies in individual centers [3]. In addition, most studies have examined only daily counts of ED visits, even though information on time of registration in the emergency department affords the opportunity of examining associations on a shorter time scale. In this study, we set out to examine associations of a comprehensive array of respiratory and cardiac conditions with the full suite of conventional air pollutants in a large multi-center study, using both daily and 3 hour average measures of exposure and response.

METHODS

Air pollution data were obtained from the National Air Pollution Surveillance (NAPS) system, and weather data from Environment Canada's weather archive. We obtained data on carbon monoxide (CO), nitrogen dioxide (NO₂), ozone (O₃), sulfur dioxide (SO₂), and particulate matter of median aerodynamic diameter less than 10 and 2.5 microns (PM₁₀, PM_{2.5} respectively). CO, NO₂, O₃ and SO₂ were measured using "reference methods" or "equivalent methods" as designated by the United States Environmental Protection Agency. CO was measured using non-dispersive infrared spectrometry, NO₂ using chemiluminescence, O₃ using chemiluminescence/ultraviolet photometry and SO₂ using coulometry/ultraviolet fluorescence. PM_{2.5} and PM₁₀ were measured using tapered element oscillating microbalance instruments. When data were available from more than one monitoring site in a city, they were averaged as a measure of average community exposure in each city.

Administrative ED visit data are not routinely captured in electronic form from all provinces. Two provinces in this study (Alberta and Ontario) collect some administrative data; however, most ED data are housed in individual institutions [4]. We therefore obtained data from individual institutions which had created their own electronic databases as follows: Montreal (Sir Mortimer B. Davis Jewish General Hospital); Ottawa (Ottawa Civic Hospital); Edmonton (Royal Alexandra Hospital, Stollery Children's Health Centre, Sturgeon Community Hospital and Health Centre, University of Alberta Hospital, Grey Nuns Community Hospital and Health Centre, Misericordia Community Hospital and Health Centre); Saint John (Saint John Regional Hospital, St. Joseph's

Hospital); Halifax (Queen Elizabeth II Health Centre); Toronto (St. Michael's Hospital, Sunnybrook Hospital); and Vancouver (St. Paul's Hospital). There is no children's hospital in Saint John, so all visits by children were included in these data. In the case of Edmonton, data from the children's hospital were included in our data set. The other centers have children's hospitals; however, they were not included in our data.

Emergency visit data were coded for the discharge diagnosis using ICD-9 or 10 by medical records staff at each institution. In addition, data pertaining to demographics (e.g., age and sex) and administrative details (e.g., date and time of visit) were obtained for every ED visit. We defined diagnosis groups as follows (ICD-9 and –10 respectively): angina/ myocardial infarction (410-414; I20-I22, I24-I25); dysrhythmia/ conduction disturbance (426, 427; I44-I49); heart failure (428; I50); respiratory infection (464, 466, 480-487; J05.0, J10-J16, J18, J20-J21); asthma (493; J45); and chronic obstructive pulmonary disease (490-492, 494-496; J40-J44, J47, J67). A control series (anemia, epilepsy, selected nerve disorders (eg. trigeminal neuralgia), selected muscle disorders (eg. muscular dystrophy), selected gastrointestinal disorders (eg. appendicitis), chronic liver disease and cirrhosis, selected gall bladder and intestinal conditions, calculus of lower urinary tract, and hyperplasia of the prostate) was also constructed (280-281.9,345, 350-356, 358-359.5,530-534, 540-543, 560-569, 571,572, 574-578, 594, 600; D50-D53, G40-G41, G50-G52, G54, G57, G58, G60, G70, G71.0-G71.3, G72.1-G72.3, K20-K22, K25-K28, K35-K38, K56-K59, K60-K63, K65-K66, K70, K72.1, K72.9, K73, K74, K75.0, K75.1, K76.6, K76.7, K80-K83, K85-K86, K92.0-K92.2, K92.8, K92.9, N21, N40). ICD-10 nomenclature was introduced at various times at different institutions. The study was approved by the individual research ethics boards at each participating

institution and the data were transferred to the Health Canada team following de-identification. No patient contact was made and patients could not be traced.

The daily number of ED visits was modelled as a function of the exponential of air pollution concentrations using generalized linear models in S Plus. The error distribution was specified as quasi-Poisson, where the variance is proportional to the expected response, accommodating over/under dispersion relative to Poisson variation. Natural spline functions of time were employed to adjust for seasonal cycles in air pollution and health outcomes. The number of knots of the cubic polynomials that comprise the splines specify the degree of non-linearity in the spline functions. In order to determine optimal temporal adjustments, natural spline smooths were constructed based on knots placed every n weeks for $n=1$ to 52. Optimality of smoothing was judged objectively based on minimization of the Akaike Information Criterion and maximization of the Bartlett's test p-value. Sensitivity analyses were conducted where associations were statistically significant, employing knots at every 3, 5 and 17 weeks (approximately 17, 10 and 3 knots per year). Indicator functions were also created for day of week and major Canadian holidays (New Year's Day, Good Friday, Victoria Day, St. Jean Baptiste Day, Canada Day, Civic Holiday, Labour Day, Thanksgiving Day, and Christmas Day). In our analysis of sub-daily effects (see below), we also included indicator variables for time of day (eight 3-hour periods per day).

In order to parsimoniously adjust for effects of weather, we chose temperature and relative humidity as the primary meteorological factors associated with visits. Mean daily temperature and relative humidity with a lag of 0, 1 and 2 days were forced into the

model using natural spline functions with 4 degrees of freedom for each lag in order to capture potential non-linearity of the exposure-response.

After making city-specific adjustments for temporal cycles and weather as described above, air pollution terms were added. We examined effects on 2 different time scales. In the conventional analysis, we used single day lags of 0-2 days of the daily average for all pollutants. We also examined effects within a day, by considering 3 hour averages of visits versus 3 hour average pollutant concentrations (e.g. 12 a.m.-3 a.m., 3 a.m.-6 a.m., 6 a.m.- 9 a.m. etc.), lagged up to 12 hours before the time of presentation to the ED. For each lag, regression parameter estimates for each pollutant were pooled among the centers using a fixed or random effects model, depending on the occurrence of statistically significant heterogeneity among effect estimates [5]. Pooled estimates are essentially a weighted average, with results from each city assigned a weight inversely proportional to the sum of within and between study variance. Thus, cities with more precise estimates are weighted more heavily in the pooled estimate. It has been observed that multi-city estimates are more stable and precise than those based on individual cities [3]. Finally, we calculated the percentage change in visits associated with a change in the pollutant concentration equivalent to the average among all cities.

RESULTS

While data for the pollutant gases were available for essentially all days, particulate matter data were less consistently available. Average air pollution concentrations among sites (Table 1) were not widely divergent, with the exception of NO₂ concentrations which tended to be lower in Saint John, and SO₂ concentrations, which were higher in Halifax and Saint John. Edmonton, for which visit data were available for multiple institutions, accounted for nearly 70% of all visits (Table 2). Since visit data were only available from selected institutions in each city, visit counts are not proportional to population. Respiratory infections, myocardial infarction/ angina and asthma were responsible for the largest number of visits in all centers, while COPD, heart failure and dysrhythmia/ conduction disturbance accounted for the fewest. Visits for heart failure exhibited the largest percentage of visits by individuals 65 years or older (76-90% among the centers), followed by COPD (49-76%), and myocardial infarction/angina (53-68%). Visits for asthma were associated with the smallest percentage of individuals 65 or older (4-20%). Visits by children (under 16 years) for asthma and respiratory infections were most frequent in Saint John (44 and 40% respectively) where there is no children's hospital, and Edmonton (41 and 47%) where data from the children's hospital were available, compared to the other sites (0-8% and 0-6%).

Associations between daily average air pollution concentrations, lagged 0-2 days, and cardiac and respiratory visits are shown in Tables 3 and 4. Several positive, statistically significant associations were observed with visits for cardiac conditions, while two negative, statistically significant associations were observed. Associations over multiple lags and/or conditions were observed for CO (lag 0 for both

angina/myocardial infarction and heart failure), NO₂ (lag 0 and 1 for angina /myocardial infarction and lag 0 for heart failure) and particulate matter with myocardial infarction/angina (PM₁₀ lag 1) and heart failure (PM₁₀ and PM_{2.5} lag 0) as well as SO₂ with myocardial infarction/angina (lag 0 and 1). Few statistically significant associations (positive or negative) were observed for respiratory conditions. However, ozone was associated with both asthma and COPD visits (lag 2). Associations of eight hour maximum ozone concentration with asthma and COPD visits were similar or smaller in magnitude than associations based on 24-hour average concentration (results not shown). No associations were observed with control conditions.

Effects of CO and NO₂ on myocardial infarction/angina and heart failure by city are shown in Figures 1 and 2. Pooled estimates are shown both with and without Edmonton in order to examine the consistency of effects relative to the centre with the largest sample size and hence most precise estimates assigned the greatest weight in calculating the pooled estimate. In the case of myocardial infarction/angina, effect sizes in Edmonton and Ottawa were similar in magnitude, while effect sizes in other centers were more variable and less precise. With respect to heart failure, effects in Edmonton and Montreal were closer in magnitude and those elsewhere tended to be more variable and less precise. The effects of ozone on asthma and COPD visits by city are shown in Figure 3. Effect size estimates were consistently less precise in centers other than Edmonton. With the exception of the association between NO₂ and myocardial infarction/angina visits, pooled estimates were generally similar in magnitude when Edmonton was excluded, and somewhat less precise. This indicates both that the pooled estimate is not singularly reflecting the estimate from Edmonton and that the pooled estimate can be

statistically significant even when effects in individual centers are not, because of the collective power of the pooled sample. There was no evidence of heterogeneity among centers ($p > 0.05$ for Q-statistic).

Associations of CO and NO₂ with myocardial infarction/angina and heart failure tended to be of larger magnitude during the warm season. CO (lag 0) was associated with a 5.2% increase in myocardial infarction/angina visits (95% CI 0.4-10.2) per 0.7 ppm, and a 6.8% increase in heart failure visits (95% CI -0.8-14.9), which were approximately twice the magnitude of the effects seen over the whole year. Similarly, NO₂ was associated with a 4.0% increase in myocardial infarction/angina visits (95% CI -0.5-8.8) per 18.4 ppb, and a 7.2% increase in heart failure visits (95% CI 0.5-14.4). These effects were approximately 50% larger than those seen over the whole year. Effects of PM_{2.5} and SO₂ on myocardial infarction/angina were weaker in the individual seasons compared to the analysis over the entire year. The magnitude of the association between ozone and COPD visits was nearly twice as large during the warm season as over the whole year (6.2% increase in visits, 95% CI, 0.1-12.7, per 18.4 ppb) and the associations of PM₁₀ and PM_{2.5} with asthma visits were respectively nearly three- and over fourfold larger (14.4% increase in visits, 95% CI, 0.2-30.7, per 20.6 $\mu\text{g}/\text{m}^3$ PM₁₀ and 7.6% increase in visits, 95% CI, 5.1-10.1, per 8.2 $\mu\text{g}/\text{m}^3$ PM_{2.5}). No consistent associations were observed between any pollutants and cardiac or respiratory visits during the winter. The magnitude of effects was generally not sensitive to the number of knots in the natural spline, but associations of SO₂ with MI/angina and of CO and NO₂ with heart failure were only statistically significant in the base analysis where the number of knots in the natural spline was optimized separately in each city (Figure 4). The effect of ozone on asthma

was much larger with knots at every 17 weeks. In a two pollutant model of MI/angina visits, including CO and NO₂, the effects of both pollutants were reduced in magnitude and statistical significance. At their mean concentrations, CO was associated with a 0.16% increase in visits ($p = 0.68$) and NO₂ was associated with a 1.18% increase in visits ($p = 0.55$). There was no evidence of consistent associations between any pollutant and cardiac or respiratory visits on sub-daily time scales (results not presented but available upon request).

DISCUSSION

We have identified significant effects of air quality on acute cardiac and respiratory presentations in a large, multi-center Canadian study of air pollution and ED visits. Unique features of this study include application of a consistent methodology to, and pooling of results from multiple diverse sites, each with over 10,000 visits, with a total of nearly 400,000 visits; examination of a comprehensive suite of cardiac and respiratory conditions; evaluation of effects over multiple time scales; and assessment of effects at generally lower pollutant concentrations than reported elsewhere. With the exception of a study in Atlanta of approximately 1 million visits for cardiovascular and respiratory disease [6, 7], many other recent studies have been based on fewer than 10,000 visits, and have examined single conditions or were restricted to specific age groups or seasons.

Compared to other recent studies in Canada of the same sites, our present results confirm our earlier findings in Saint John of associations between ozone and asthma and COPD visits [8, 9]. In contrast, we found no consistent associations in the current study

with respiratory infections, despite comprising the largest single diagnostic category. We found significant associations, however, between cardiac visits and CO and NO₂, which were not detected in Saint John. As a small city, Saint John does not have large traffic volumes, so it is characterized by low ambient concentrations of these pollutants.

Villeneuve et al. [10] also reported associations of CO and NO₂ with ED visits for stroke and transient ischemic attacks. Conversely, Saint John has relatively high concentrations of SO₂. There was some evidence of an association between SO₂ and angina/ myocardial infarction in the current analysis, but not with any of the respiratory diagnoses.

Elsewhere, several recent studies have evaluated associations between air pollution and ED visits for respiratory disease. CO exhibited an association with COPD [7, 11] and upper respiratory infection (URI) visits [7]. Associations were observed between NO₂ and visits for asthma [12, 13], COPD [7], URI [7] and lower respiratory disease [14]. Ozone was associated with respiratory visits in several studies, including visits for URI [7], asthma [13, 15, 16], and COPD [11]. Associations with particulate matter were observed for asthma [13, 15] as well as URI and pneumonia [7]. Finally, SO₂ exhibited associations with asthma visits [13, 15, 16]. A study of CO, PM₁, PM_{2.5} and PM₁₀ relative to respiratory visits in Spokane, revealed generally weak and inconsistent associations [17]. The authors also examined associations with hospital admissions and mortality and again results were inconsistent. Fewer studies have examined associations between air pollution and visits for cardiovascular disease. In a study in Atlanta, CO, NO₂ and PM_{2.5} (and specific components) were associated with visits for cardiovascular disease [6]. CO, NO₂, SO₂ and PM_{2.5} were associated with visits for cardiovascular

disease in those 65 years of age or older in Sydney [18]. Only CO was associated with visits for angina or myocardial infarction in a study in Sao Paulo [19].

We observed associations between daily average air pollution concentrations and emergency department visits with lags of zero to two days, but not for three hour average concentrations lagged up to 12 hours within the same day. We previously reported that approximately 30% and 70% respectively of those visiting the emergency department with respiratory or cardiac conditions did so within less than one day of developing the symptoms most responsible for their visit [2]. This is consistent with our current findings of predominant effects of CO and NO₂ on cardiac conditions at lag 0, and of O₃ on respiratory conditions at lag 2. In contrast, Villeneuve et al. [20] reported significant associations between 6 hour average weather variables such as thunderstorm activity, and visits by children for asthma.

Somewhat larger effects were observed for visits for heart failure compared to myocardial infarction. Individuals with heart failure have been identified as one group at particular risk of air pollution related mortality [21]. Of respiratory conditions, effects of ozone were larger for COPD compared to asthma visits, particularly in the warm season. Individuals with COPD could be considered to have a similar relative degree of frailty compared to those with asthma, as those with heart failure compared to myocardial infarction/ angina. Indeed, in all centers in the current analysis, those making visits for COPD were, not surprisingly, older than those with asthma, just as those presenting with heart failure were older than those with myocardial infarction/ angina.

We also observed larger effects in the warm season, which is consistent with knowledge of time-activity patterns [22]. These data indicate that individuals spend more

time outdoors during warm weather, increasing their exposure to outdoor air pollution and by consequence reducing exposure misclassification of exposures derived from fixed site monitoring stations. This lends additional credibility to the existence of a causal association between exposure and response.

While the pathophysiology of the impacts of air pollution on the respiratory system has been well documented over many years, particularly in relation to ozone and particulate matter, mechanisms of action on the cardiovascular system have received more recent attention. These include effects on arterial vasoconstriction [23], heart rate variability [24, 25], blood clotting [26] and formation of atherosclerotic plaques [27]. Several studies have documented an increased frequency of discharges of implantable defibrillators related to short-term increases in exposure to particulate matter, CO and NO₂ [28, 29]. Peters et al. [30] recently reported an association between exposure to traffic and onset of myocardial infarction.

Limitations

We relied on diagnostic information provided in administrative data records, which would be expected to be associated with a degree of misclassification. However, in an earlier evaluation of emergency department diagnostic classification for respiratory and cardiac conditions, we observed substantial or better interobserver agreement in diagnostic classification ($\kappa = 0.69$ to 0.84 for the categories considered here), respiratory infections notwithstanding ($\kappa = 0.53$), and there was no evidence of diagnostic bias in relation to daily air pollution level [31]. Although compared to many recent studies, we had the advantage of a relatively large sample size, our ability to detect effects of

particulate matter may have been constrained by the limited availability of data in some centers. Applying a consistent methodology to multiple sites is also advantageous, but one center (Edmonton) had a disproportionately large sample. Nonetheless, sample sizes from other sites, or for specific conditions pooled over multiple sites were still larger than most recent studies of ED visits, and pooled estimates were generally similar, although less precise, when Edmonton was excluded. Since, with the exception of Edmonton and Saint John, data were available only for selected hospitals, these may not be representative of the entire population in a given city. However, in the context of Canada's publicly funded universal healthcare system, large differences between hospitals are less likely. Statistically significant heterogeneity among sites was not detected, probably because of the large within versus between site variance of effects. While we have attempted to focus on effects which exhibited consistency over multiple lags and/or diagnosis groups, we have nonetheless conducted numerous hypothesis tests, increasing the probability of false positive results. However, the number of positive, statistically significant associations with visits for cardiac conditions exceeded the number which could be expected by chance alone, and few negative and significant associations were observed.

CONCLUSIONS

We conducted a time series analysis of air pollution and cardiac and respiratory emergency department visits in seven Canadian cities, based on nearly 400,000 visits to 14 hospitals. CO and NO₂ exhibited the most consistent associations with cardiac conditions, while ozone was most consistently associated with respiratory visits. PM₁₀

and $PM_{2.5}$ were strongly associated with asthma visits during the warm season. No consistent associations were observed between cardiac or respiratory visits and three hour average pollutant concentrations lagged up to 12 hours. In addition to their consistency with other results, these results add further weight to arguments regarding the role of air pollution in contributing to adverse health events, and imply that interventions to reduce these pollutants are warranted in an effort to reduce cardio-respiratory ED visits.

LIST OF ABBREVIATIONS USED

CO-carbon monoxide

COPD-chronic obstructive pulmonary disease

ED-emergency department

ICD- international classification of diseases

MI-myocardial infarction

NAPS-national air pollution surveillance

NO₂- nitrogen dioxide

O₃- ozone

PM₁-particulate matter of median aerodynamic diameter less than 1 μm

PM_{2.5}-particulate matter of median aerodynamic diameter less than 2.5 μm

PM₁₀- particulate matter of median aerodynamic diameter less than 10 μm

ppb-parts per billion

ppm-parts per million

SO₂- sulfur dioxide

URI-upper respiratory infection

COMPETING INTERESTS: The authors declare that they have no competing interests.

AUTHORS' CONTRIBUTIONS: DMS conceived of the study, directed the statistical analysis and drafted the manuscript. MS conducted the statistical analysis. BHR participated in study design and directed data acquisition. JAL participated in study design and contributed data. All authors contributed to drafting the manuscript and read and approved the final version.

AUTHORS' INFORMATION: DMS is a public health physician in the Population Studies Division, Healthy Environments and Consumer Safety Branch of Health Canada, an adjunct professor in the Department of Epidemiology and Community Medicine at the University of Ottawa, and an affiliate scientist at the McLaughlin Center for Population Health Risk Assessment. MS is a research scientist in the Population Studies Division, Healthy Environments and Consumer Safety Branch of Health Canada. BHR is a clinical emergency physician, Professor and Research Director in the Division of Emergency Medicine, University of Alberta, and Canada Research Chair in Emergency Airway Diseases. JAL is a respirologist in the Division of Respiriology, Department of Medicine, University of Ottawa, and medical epidemiologist in the Population Studies Division, Healthy Environments and Consumer Safety Branch of Health Canada

ACKNOWLEDGMENTS: Drs. Robert Beveridge, Eric Grafstein, Simon Kingsley, Michael Schull, Douglas Sinclair and Bernard Unger provided emergency department visit data. Serge Beaulieu of Atlantic Health Sciences Corporation, Stacey Ackroyd of

Dalhousie University, Chris Houston from Information Services, Capital Health and Virginia Willis, from the Emergency Medicine Research Group (EMeRG), assisted with data access. The study was funded by Health Canada and Environment Canada.

REFERENCES

1. Stieb DM, Judek S, Burnett RT: **Meta-analysis of time-series studies of air pollution and mortality: effects of gases and particles and the influence of cause of death, age, and season.** *J Air Waste Manag Assoc* 2002, **52**:470-84.
2. Stieb DM., Beveridge RC, Smith-Doiron M, Burnett RT, Judek S, Dales RE, Anis AH: **Beyond administrative data: characterizing cardiorespiratory disease episodes among patients visiting the emergency department.** *Can J Public Health* 2000, **91**:107-112.
3. Dominici F, Samet JM, Zeger SL: **Combining evidence on air pollution and daily mortality from the twenty largest US cities: A hierarchical modeling strategy (with discussion).** *R Stat Soc* 2000, **163**:263-302.
4. Rowe BH, Bond K, Ospina M, Blitz S, Schull MJ, Sinclair D, Bullard M: **Data collection on patients in emergency departments in Canada.** *Can J Emerg Med* 2006, **8**: 417-424.
5. DerSimonian R, Laird N: **Meta-analysis in clinical trials.** *Control Clin Trials* 1986, **7**:177-88.
6. Metzger KB, Tolbert PE, Klein M, Peel JL, Flanders WD, Todd K, Mulholland JA, Ryan PB, Frumkin H: **Ambient air pollution and cardiovascular emergency department visits.** *Epidemiology* 2004, **15**:46-56.
7. Peel, JL, Tolbert PE, Klein M, Metzger KB, Flanders WD, Todd K, Mulholland JA, Ryan PB, Frumkin H: **Ambient air pollution and respiratory emergency department visits.** *Epidemiology* 2005, **16**:164-174.

8. Stieb DM, Burnett RT, Beveridge RC, Brook JR: **Ozone and asthma emergency department visits in Saint John, New Brunswick, Canada.** *Environ Health Perspect* 1996, **104**:1354-1360.
9. Stieb DM, Beveridge RC, Brook JR, Smith-Doiron M, Burnett RT, Dales RE, Beaulieu S, Judek S, Mamedov A: **Air pollution, aeroallergens and cardiorespiratory emergency department visits in Saint John, Canada.** *J Expo Anal Environ Epidemiol* 2000, **10**:461-477.
10. Villeneuve PJ, Chen L, Stieb D, Rowe BH: **Associations between outdoor air pollution and emergency department visits for stroke in Edmonton, Canada.** *Eur J Epidemiol* 2006, **21**:689-700.
11. Tenias JM, Ballester F, Perez-Hoyos S, Rivera ML: **Air pollution and hospital emergency room admissions for chronic obstructive pulmonary disease in Valencia, Spain.** *Arch Environ Health* 2002, **57**:41-47.
12. Erbas B, Kelly AM, Physick B, Code C, Edwards M: **Air pollution and childhood asthma emergency hospital admissions: estimating intra-city regional variations.** *Int J Environ Health Res* 2005, **15**:11-20.
13. Galan I, Tobias A, Banegas JR, Aranguiz E: **Short-term effects of air pollution on daily asthma emergency room admissions.** *Eur Respir J* 2003, **22**:802-808.
14. Farhat SC, Paulo RL, Shimoda TM, Conceicao GM, Lin CA, Braga AL, Warth MP, Saldiva PH: **Effect of air pollution on pediatric respiratory emergency room visits and hospital admissions.** *Braz J Med Biol Res* 2005, **38**:227-235.
15. Jaffe DH, Singer ME, Rimm AA: **Air pollution and emergency department visits for asthma among Ohio Medicaid recipients, 1991-1996.** *Environ Res* 2003, **91**:21-28.

16. Wilson AM, Wake CP, Kelly T, Salloway JC: **Air pollution, weather, and respiratory emergency room visits in two northern New England cities: an ecological time-series study.** *Environ Res* 2005, **97**:312-321.
17. Slaughter JC, Kim E, Sheppard L, Sullivan JH, Larson TV, Claiborn C: **Association between particulate matter and emergency room visits, hospital admissions and mortality in Spokane, Washington.** *J Expo Anal Environ Epidemiol.* 2005, **15**:153-159.
18. Jalaludin B, Morgan G, Lincoln D, Sheppard V, Simpson R, Corbett S: **Associations between ambient air pollution and daily emergency department attendances for cardiovascular disease in the elderly (65+ years), Sydney, Australia.** *J Expo Anal Environ Epidemiol* 2006, **16**:225-237
19. Lin CA, Amador Pereira LA, de Souza Conceicao GM, Kishi HS, Milani R Jr, Ferreira Braga AL, Nascimento Saldiva PH: **Association between air pollution and ischemic cardiovascular emergency room visits.** *Environ Res* 2003, **92**:57-63.
20. Villeneuve PJ, Leech J, Bourque D: **Frequency of emergency room visits for childhood asthma in Ottawa, Canada: the role of weather.** *Int J Biometeorol* 2005, **50**:48-56.
21. Goldberg MS, Burnett RT, Valois MF, Flegel K, Bailar JC 3rd, Brook J, Vincent R, Radon K: **Associations between ambient air pollution and daily mortality among persons with congestive heart failure.** *Environ Res* 2003, **91**:8-20.
22. Leech JA, Nelson WC, Burnett RT, Aaron S, Raizenne ME: **It's about time: a comparison of Canadian and American time-activity patterns.** *J Expo Anal Environ Epidemiol* 2002, **12**:427-432.

23. Brook RD, Brook JR, Urch B, Vincent R, Rajagopalan S, Silverman F: **Inhalation of fine particulate air pollution and ozone causes acute arterial vasoconstriction in healthy adults.** *Circulation* 2002, **105**:1534-1536.
24. Pope CA 3rd, Hansen ML, Long RW, Nielsen KR, Eatough NL, Wilson WE, Eatough DJ: **Ambient particulate air pollution, heart rate variability, and blood markers of inflammation in a panel of elderly subjects.** *Environ Health Perspect* 2004, **112**:339-345.
25. Wheeler A, Zanobetti A, Gold DR, Schwartz J, Stone P, Suh HH: **The relationship between ambient air pollution and heart rate variability differs for individuals with heart and pulmonary disease.** *Environ Health Perspect* 2006, **114**:560-566.
26. Peters A, Doring A, Wichmann H-E, Koenig W: **Increased plasma viscosity during an air pollution episode: a link to mortality?** *Lancet* 1997, **349**:1582-1587.
27. Künzli N, Jerrett M, Mack WJ, Beckerman B, LaBree L, Gilliland F, Thomas D, Peters J, Hodis HN: **Ambient air pollution and atherosclerosis in Los Angeles.** *Environ Health Perspect* 2005, **113**:201-206.
28. Dockery DW, Luttmann-Gibson H, Rich DQ, Link MS, Mittleman MA, Gold DR, Koutrakis P, Schwartz JD, Verrier RL: **Association of air pollution with increased incidence of ventricular tachyarrhythmias recorded by implanted cardioverter defibrillators.** *Environ Health Perspect* 2005, **113**:670-674.
29. Peters A, Liu E, Verrier RL, Schwartz J, Gold DR, Mittleman M, Baliff J, Oh JA, Allen G, Monahan K, Dockery DW: **Air pollution and incidence of cardiac arrhythmia.** *Epidemiology* 2000, **11**:11-17.

30. Peters A, von Klot S, Heier M, Trentinaglia I, Hörmann A, Wichmann H-E, Löwel H: **Exposure to traffic and the onset of myocardial infarction.** *N Engl J Med* 2004, **351**:1721-1730.
31. Stieb DM, Beveridge RC, Rowe BH, Walter SD, Judek S: **Assessing diagnostic classification in an emergency department: implications for daily time series studies of air pollution.** *Am J Epidemiol* 1998, **148**:666-670.

Figure titles (legends)

Figure 1. Percent increase in emergency department visits for angina/myocardial infarction by center (Point estimates and 95% confidence intervals are shown, for CO (panel A) and NO₂ (panel B)).

Figure 2. Percent increase in emergency department visits for heart failure by center. (Point estimates and 95% confidence intervals are shown, for CO (panel A) and NO₂ (panel B)).

Figure 3. Percent increase in respiratory emergency department visits by center. (Point estimates and 95% confidence intervals are shown for asthma (panel A) and COPD (panel B)).

Figure 4. Percent increase in emergency department visits for selected pollutants, diagnostic groups per mean pollutant concentration. (Point estimates and 95% confidence intervals are shown by number of knots in natural spline functions of time; from left to right, base analysis where number of knots is optimized separately in each city, and knots every 3, 5 and 17 weeks)

Table 1. Average 24 hour pollutant concentrations by city

Centre (dates)	CO (ppm)	NO ₂ (ppb)	O ₃ (ppb)	PM ₁₀ (µg/m ³)	PM _{2.5} (µg/m ³)	SO ₂ (ppb)	Percent of days with data	
							PM ₁₀	PM _{2.5}
Montreal (1/97-12/02)	0.5	19.4	18.3	25.8	8.6	4.8	49.8	88.5
Ottawa (4/92-12/00)	0.9	18.8	17.5	20.1	6.7	3.9	11.7	31.0
Edmonton (4/92-3/02)	0.7	21.9	18.6	22.6	8.5	2.6	77.0	31.2
Saint John (7/92-3/96)	0.5	9.3	20.1	NA*	NA	7.7	0.0	0.0
Halifax (1/99-12/02)	0.5	17.5	22.1	NA	9.8	10.0	0.0	33.2
Toronto (4/99-3/03)	1.0	22.7	22.0	20.7	9.1	4.2	58.1	100.0
Vancouver (6/99-12/02)	0.6	18.7	10.3	13.6	6.8	2.6	100.0	71.2
Average	0.7	18.4	18.4	20.6	8.2	5.1		

*N/A – not available

Table 2. Frequency of visits by center and diagnosis group

Center	Myocardial Infarction/ Angina	Heart Failure	Dysrhythmia/ Conduction Disturbance	Asthma	Chronic Obstructive Pulmonary Disease	Respiratory Infection	Total
Montreal	4,978	3,833	3,943	2,925	2,394	4,761	22,834
Ottawa	12,156	5,115	6,912	5,873	4,764	8,222	43,042
Edmonton	35,207	17,115	26,813	62,563	26,527	90,509	258,734
Saint John	2,435	1,312	1,096	4,771	1,761	8,446	19,821
Halifax	2,834	1,661	1,643	2,815	1,978	5,321	16,252
Toronto	3,194	2,035	2,897	2,652	1,827	3,563	16,168
Vancouver	2,380	1,242	1,856	1,964	1,240	4,323	13,005
Total	63,184	32,313	45,160	83,563	40,491	125,145	389,856

Table 3. Percent increase in cardiac visits (95% confidence interval) for specified change in pollutant concentration.

Pollutant	Lag	Mean	Angina/ Myocardial Infarction	Heart Failure	Dysrhythmia
CO	0	0.7	2.1 (0.0, 4.2)*	3.8 (0.7, 6.9)	-2.7 (-5.8, 0.5)
	1	ppm	1.6 (-0.5, 3.7)	2.8 (-2.1, 8.0)	-4.0 (-10.1, 2.5)
	2		-0.2 (-2.2, 1.8)	1.7 (-2.7, 6.2)	-2.5 (-8.4, 3.8)
NO ₂	0	18.4	2.6 (0.2, 5.0)	4.7 (1.2, 8.4)	-1.3 (-4.1, 1.5)
	1	ppb	2.7 (0.2, 5.3)	2.8 (-1.3, 7.1)	-0.9 (-3.8, 2.1)
	2		0.7 (-1.6, 3.1)	1.9 (-3.1, 7.1)	0.3 (-2.6, 3.3)
O ₃	0	18.4	-3.4 (-7.8, 1.3)	-1.2 (-6.5, 4.5)	-1.7 (-6.6, 3.4)
	1	ppb	<i>-3.0 (-5.9, 0.0)†</i>	-0.5 (-4.4, 3.6)	2.0 (-3.7, 7.9)
	2		-0.3 (-4.0, 3.5)	2.6 (-1.4, 6.7)	1.5 (-4.7, 8.2)
PM ₁₀	0	20.6	0.0 (-2.1, 2.2)	9.3 (0.8, 18.4)	-0.1 (-5.9, 6.1)
	1	µg/m ³	1.9 (-0.3, 4.1)	0.4 (-2.8, 3.6)	-0.9 (-3.4, 1.6)
	2		-0.3 (-2.5, 1.8)	-0.1 (-3.2, 3.0)	2.1 (-4.0, 8.5)
PM _{2.5}	0	8.2	2.0 (-1.3, 5.5)	6.5 (0.1, 13.4)	-0.8 (-3.1, 1.5)
	1	µg/m ³	1.1 (-2.0, 4.3)	-0.1 (-2.8, 2.7)	0.0 (-2.3, 2.3)
	2		0.9 (-1.0, 2.8)	-0.4 (-3.0, 2.3)	0.0 (-2.3, 2.3)
SO ₂	0	5.1	1.7 (-0.2, 3.5)	0.9 (-2.8, 4.7)	-0.7 (-3.1, 1.7)
	1	ppb	2.1 (0.2, 4.0)	1.5 (-1.0, 4.1)	0.4 (-3.3, 4.3)
	2		0.0 (-2.3, 2.4)	0.8 (-1.6, 3.3)	-2.6 (-4.8, -0.3)

*Positive, statistically significant associations (are shown in bold. †Negative, statistically significant associations are shown in italics.

Table 4. Percent increase in respiratory visits (95% confidence interval) for specified change in pollutant concentration.

Pollutant	Lag	Mean	Asthma	Chronic Obstructive Pulmonary Disease	Respiratory Infection
CO	0	0.7	-1.8 (-5.8, 2.2)	2.2 (-3.4, 8.1)	-0.8 (-3.2, 1.7)
	1	ppm	-0.3 (-2.3, 1.8)	<i>-3.3 (-6.1, -0.4)†</i>	0.7 (-3.0, 4.5)
	2		0.6 (-1.5, 2.7)	-2.3 (-6.1, 1.6)	2.4 (-1.7, 6.7)
NO ₂	0	18.4	-0.4 (-4.4, 3.9)	0.1 (-5.6, 6.2)	-0.9 (-2.9, 1.1)
	1	ppb	-1.2 (-4.6, 2.3)	<i>-3.4 (-6.6, -0.1)</i>	0.7 (-3.7, 5.3)
	2		0.0 (-2.4, 2.5)	-4.8 (-11.5, 2.5)	0.6 (-1.4, 2.6)
O ₃	0	18.4	0.1 (-3.8, 4.2)	-2.2 (-5.9, 1.7)	-0.6 (-3.1, 1.9)
	1	ppb	4.3 (-1.3, 10.2)	2.2 (-1.7, 6.3)	0.1 (-2.3, 2.5)
	2		3.2 (0.3, 6.2)*	3.7 (-0.5, 7.9)	-0.1 (-3.0, 2.9)
PM ₁₀	0	20.6	5.3 (-2.2, 13.4)	-0.6 (-3.3, 2.2)	0.0 (-1.8, 1.8)
	1	µg/m ³	0.9 (-1.1, 3.0)	-1.7 (-7.6, 4.5)	-0.9 (-2.7, 0.9)
	2		1.1 (-2.4, 4.8)	-5.3 (-11.4, 1.2)	-1.2 (-3.0, 0.6)
PM _{2.5}	0	8.2	1.7 (-2.5, 6.1)	-1.8 (-6.1, 2.7)	0.6 (-1.4, 2.7)
	1	µg/m ³	0.9 (-1.1, 3.0)	-0.9 (-3.6, 1.8)	0.6 (-1.1, 2.4)
	2		0.4 (-1.6, 2.4)	-0.2 (-5.0, 4.8)	-0.2 (-1.9, 1.5)
SO ₂	0	5.1	-1.1 (-2.9, 0.7)	-1.9 (-4.3, 0.6)	-0.6 (-2.1, 0.9)
	1	ppb	-1.8 (-4.2, 0.7)	-1.4 (-4.4, 1.8)	-0.2 (-1.7, 1.3)
	2		-1.1 (-2.8, 0.7)	-3.0 (-6.7, 0.9)	0.6 (-1.3, 2.6)

*Positive, statistically significant associations are shown in bold. †Negative, statistically significant associations are shown in italics.

Angina/Myocardial Infarction






Figure 1

Heart Failure

A



B





Figure 2

A) Asthma



B) COPD




Figure 3




Figure 4