

**Aryl Hydrocarbon Receptor (AhR) Agonists Suppress  
Interleukin-6 Expression by Bone Marrow Stromal Cells**

Brenda A. Jensen, Rebecca J. Leeman, Jennifer J. Schlezinger and David H. Sherr\*

Department of Environmental Health, Boston University School of Public Health, 715 Albany Street,  
Boston, MA 02118

\*Correspondence to:

David H. Sherr, Department of Environmental Health, Boston University School of Public Health, 715  
Albany Street (R-408), Boston, Massachusetts, 02118, Phone: 617-638-6464; FAX: 617-638-6463; E-  
mail: dsherr@bu.edu



## ABSTRACT

**Background:** Bone marrow stromal cells produce cytokines required for the normal growth and development of all eight hematopoietic cell lineages. Aberrant cytokine production by stromal cells contributes to blood cell dyscrasias. Consequently, factors that alter stromal cell cytokine production may significantly compromise the development of normal blood cells. We have shown that environmental chemicals, such as aromatic hydrocarbon receptor (AhR) agonists, suppress B lymphopoiesis by modulating bone marrow stromal cell function. Here, we extend these studies to evaluate the potential for two prototypic AhR agonists, 7,12-dimethylbenz[*a*]anthracene (DMBA) and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD), to alter stromal cell cytokine responses.

**Methods:** Bone marrow stromal cells were treated with AhR agonists and bacterial lipopolysaccharide (LPS) to mimic innate cytokine responses to bacterial challenges and to study the effects of AhR ligands on those responses. Steady state cytokine RNA levels were screened by RNase protection assays (RPA) and quantified by real-time PCR. Cytokine (IL-6) protein production was measured by ELISA. ~~and transcription factor (NF- $\kappa$ B) activity by~~ EMSAs were used to study IL-6 transcriptional regulation.

**Results:** RPAs indicated that AhR<sup>+</sup> bone marrow stromal cells up-regulated genes encoding IL-6, LIF, G-CSF, and GM-CSF in response to LPS, presumably through activation of Toll-like receptor 4. Pre-treatment with low doses of DMBA or TCDD selectively abrogated *IL-6* gene induction but had no effect on the other up-regulated genes. Real-time-PCR indicated a significant inhibition of *IL-6* mRNA by AhR ligands within 1 hour of LPS challenge which was reflected in a profound down-regulation of IL-6 protein induction, with DMBA and TCDD suppressing IL-6 levels as much as 65% and 88%, respectively. This potent inhibitory effect persisted for at least 72 hours. EMSAs measuring NF- $\kappa$ B binding to IL-6 promoter sequences, an event known to induce IL-6 transcription, indicated a

significant decrease in the LPS-mediated induction of DNA-binding RelA/p50 and c-Rel/p50 heterodimers in the presence of DMBA.

**Conclusions:** Common environmental AhR agonists can suppress an innate response to [a model](#) bacterial challenge through down-regulation of IL-6, a cytokine critical to the growth of several hematopoietic cell subsets, including early B cells. This suppression occurs at least at the level of *IL-6* gene transcription and may be regulated by NF- $\kappa$ B.

## INTRODUCTION

Bone marrow stromal cells support the growth and development of all eight hematopoietic cell lineages through cell-cell contact and the production of soluble cytokines (1, 2). Although this process generally is ~~generally~~ well-regulated by both adhesion molecules and receptor/ligand pairs, aberrant stromal cell-blood cell interactions have been documented and are associated with a variety of diseases that involve abnormal growth and development of blood cells (3-5).

Perhaps the best studied stromal cell-blood cell interaction is that which occurs between bone marrow stroma and developing B lymphocytes (6-9). Of particular interest in these studies is the contribution of stromal cell-derived IL-6 to B lymphopoiesis. Originally described as a tumor-derived growth factor (10), IL-6 now is known to be required for the growth of normal bone marrow progenitor B cells (11), for terminal differentiation of surface immunoglobulin-bearing B cells (12, 13), and for the relatively long-term survival of bone marrow plasma cells (13-15).

Aberrant IL-6 expression has been associated with autoimmune diseases, including, but not limited to vitiligo (16), ~~S~~systemic ~~L~~upus ~~E~~rythematosus (SLE) (17, 18), rheumatoid arthritis (19), and multiple sclerosis (20). Moreover, IL-6 has been recognized as an important growth and survival factor for neoplastic bone marrow plasma cells in multiple myeloma (5, 14, 21) and has been targeted with specific antibodies for myeloma therapy (22). In addition, IL-6 is emerging as an important survival and angiogenesis factor in other cancers, including basal cell carcinoma, prostate cancer, and ~~Kaposs~~'s carcinoma (23-25). These studies illustrate the importance of IL-6 regulation in normal cell function and suggest that any modulation of its expression could have important pathologic consequences.

In a series of studies, ~~w~~We and others demonstrated that exposure of bone marrow stromal cells to common environmental contaminants, such as polycyclic aromatic hydrocarbons (PAH), adversely affect their function (26-34). Specifically, the prototypic PAHs, benzo[*a*]pyrene (B[*a*]P) and 7,12-

dimethylbenz[*a*]anthracene (DMBA); induce primary or cloned stromal cells to deliver a death signal to adjacent pre- and pro/pre-B cells (28-34). Induction of this apoptosis signal in stromal cells is dependent on activation of the aryl hydrocarbon receptor (AhR), a cytosolic receptor that is converted into a transcription factor on binding of any one of a number of PAH, halogenated hydrocarbons (HAHs), or planar polychlorinated biphenyls (PCBs) (35-37). Other laboratories demonstrated that HAH, such as 2,4,7,8-tetrachloro-*p*-dioxin (TCDD), suppress the production of bone marrow-derived T cell precursors (38, 39), although the role of AhR<sup>+</sup> bone marrow stromal cells in this process was not evaluated. Interestingly, exposure to AhR ligands has been associated with increased multiple myeloma risk (40-42), suggesting a possible link between AhR activation, aberrant bone marrow stromal cell cytokine production, and plasma cell dyscrasia.

In light of the ability of AhR ligands to target bone marrow stromal cells and the importance of bone marrow stromal cells to blood cell development, we sought to determine if AhR ligands compromise production of bone marrow stromal cell cytokines such as IL-6. To this end, a well-characterized AhR<sup>+</sup> bone marrow stromal cell line (BMS2) was used to evaluate the effects of a PAH (DMBA) and an HAH (TCDD) on cytokine gene expression induced by the Toll-like receptor-4 (TLR-4) ligand, lipopolysaccharide (LPS). LPS was chosen for these studies since it up-regulates the production of several bone marrow-derived growth factors, including IL-6, in models of inflammatory cytokine responses to bacterial challenge (8, 11, 43-45). Particular attention was paid to the potential role of the promiscuous transcription factor NF- $\kappa$ B in AhR ligand effects since we and others demonstrated a physical and functional interaction between the AhR and the p65 (RelA) subunit of NF- $\kappa$ B (46, 47) and since NF- $\kappa$ B plays an important role in regulating IL-6 gene transcription (48).

## METHODS

### *Chemicals*

~~DMBA and~~ TCDD ~~was~~<sup>ere</sup> obtained from Ultra Scientific (~~North Kingstown, Hope, Rhode~~  
~~Island~~<sup>RI</sup>). ~~DMBA and~~ Staphylococcus lipopolysaccharide (LPS) ~~were~~<sup>as</sup> purchased from Sigma Chemical Co. (St. Louis, MO).

### *Cell culture and treatment*

BMS2, a bone marrow stromal cell line that supports the growth and differentiation of granulocytes, pro/pre-B, and pre-B cells (8, 34) was a gift from Dr. P. Kincade (Oklahoma Medical Research Center). BMS2 cells were maintained in DMEM (Mediatech, Washington, DC) supplemented with 5% fetal calf serum, L-glutamine (Gibco/BRL, Gaithersburg, MD), 2-mercaptoethanol (Sigma Chemical Co) and 25 µg/ml plasmocin (Invivogen, Carlsbad, CA) at 37°C and 7.5% CO<sub>2</sub>. Cells were passed twice weekly and determined to be mycoplasma-free by a PCR based protocol (Mycoplasma Detection Kit; ATCC, Manassas, VA).

BMS2 cells were plated in antibiotic-free medium for 24 hours prior to treatment with vehicle (0.01% DMSO final concentration), 1-10 µM DMBA, or 1-~~0~~ nM TCDD dissolved in DMSO. One hour later, LPS was added to a final concentration of 1-~~0~~ µg/ml. Cells were harvested at various times thereafter for RNA analysis (RNase protection assays; Real-time PCR) or nuclear protein-DNA binding analyses (EMSA), and supernatant~~s~~ were harvested for IL-6 quantitation by ELISA.

### *RNA isolation*

Culture well-adherent BMS2 cells were removed with 0.05% trypsin (Sigma), rinsed with PBS, pelleted, and frozen at -80° C until use. Total RNA was isolated using the RNeasy kit (Qiagen; Valencia, CA) or SV Total RNA System (Promega, Madison, WI). RNA was quantitated and samples visualized on an RNA gel to ensure RNA integrity.

### *Ribonuclease Protection Assay (RPA)*

The RiboQuant Multi-probe RNase Protection Assay System (BD Pharmingen, San Diego, CA) was used for RPA experiments. Riboprobes were generated with kit components and fresh [ $\alpha$ - $^{32}$ P]-UTP 3000 Ci/mmol (Perkin-Elmer, [Boston, MA](#)) and used within 2 days. Two templates (mCK4 and mCK3b) were tested to screen for changes in genes encoding ~~multiple~~ several cytokines: IL-3, IL-6, IL-7, IL-11, IFN $\gamma$ , TNF $\alpha$ , TNF- $\beta$ , LT- $\beta$ , TGF $\beta$ 1, TGF $\beta$ 2, TGF $\beta$ 3, MIF, GM-CSF, M-CSF, G-CSF, LIF, and SCF. L32 and GAPDH were also represented as internal standards. Purified probes were mixed with 7-15  $\mu$ g of total RNA and hybridized overnight. After cooling, annealed targets were treated with RNase and the resulting fragments were purified, denatured and resolved on a 4.5% sequencing gel. The gels were dried completely, exposed to film and quantified with a Molecular Dynamics Phosphor Imager (Amersham Biosciences, Sunnyvale, CA) using Imagequant software (Amersham). ~~Statistics were performed with Prism version 3.0 for Macintosh (GraphPad).~~

#### *Real-Time PCR*

IL-6 primers were designed to span exon junctions and were tested against genomic templates. PCR products were resolved on agarose gels to show minimal contribution from genomic products or primer dimers. IL-6 primer sequences were: 5'CAAGAGACTTCCATCCAGTTGCCT-3' and 5'-TTTCTCATTTCACGATTTCCCAG-3'.  $\beta$ -actin primers were 5'-GTCGTCGACAACGGCTCCGGCATGTG-3' and 5'-CATTGTAGAAGGTGTGGTGCCAGATC-3' ~~Is this a mistake?~~ [Yamaguchi, 1997 #6465]. Total RNA was reverse transcribed into cDNA using TaqMan Reverse Transcription Reagents (PE Applied Biosystems, Foster City, CA). cDNA (0.01  $\mu$ g) was used in the hot start Real-time PCR with SYBR Green PCR Master Mix (PE Applied Biosystems). Real-time PCR was carried out with the ABI Prism 7700 Sequence Detector (PE Applied Biosystems). The PCR conditions were: 95°C for 10 min, 40 cycles of 90°C for 15 s, and 60°C for 1 m, with fluorescence ~~measurements~~ readings read during each cycle. Fluorescence measurements were used to

compare transcripts between samples using the Comparative  $C_T$  Method, a method for determining relative amounts of transcripts based on an internal control when the absolute number of transcripts represented is unknown (PE Applied Biosystems). ~~Statistics were conducted using Prism version 3.0 for Macintosh (GraphPad).~~

#### *ELISA*

At each time point indicated, 500  $\mu$ l aliquots of BMS2 cell culture supernatants were drawn off and frozen at  $-20^\circ\text{C}$ . ELISAs were carried out with the OptEIA mouse IL-6 ELISA kit (BD Pharmingen) according to the manufacturer's instructions. Supernatant samples were diluted 1:50 and plated in duplicate wells of a 96 well plate. Freshly diluted rIL-6 was used to generate a standard curve on each plate analyzed. Plates were read at 450 nm using a spectrophotometric plate reader. Raw data were corrected against blank wells and converted to pg/ml using the standard curve.

#### *NF- $\kappa$ B EMSA*

BMS2 cells were treated in ~~6-well plates~~~~T75 flasks~~ with vehicle or 1  $\mu\text{M}$  DMBA. One hour later, 1  $\mu\text{g}/\text{ml}$  LPS ~~or PBS~~ was added ~~to half of~~~~to~~ the cultures, and cells ~~were~~ harvested 16 or 24 hours later. Nuclear proteins were extracted as described previously (49). For determination of NF- $\kappa$ B activation, a double stranded oligonucleotide containing the NF- $\kappa$ B binding site from the upstream regulatory element of *c-myc* (5'-GATCCAAGTCCGGGTTTTCCCAACC-3') (50) was used. The DNA probe was end-labeled using T4 polynucleotide kinase (Promega, Madison, WI) and [ $\gamma$ - $^{32}\text{P}$ ]-ATP and was purified using a Centriscin-20 column (Princeton Separations, Adelphia, NJ). EMSAs were performed as follows:  $^{32}\text{P}$ -labeled DNA ( $\sim 0.5$  ng, 50,000 cpm) and 2  $\mu\text{g}$  of nuclear protein were combined with buffer (final concentrations: 10mM Tris-HCl, pH 7.5, 1 mM EDTA, 100 mM sodium chloride, 0.5 mM magnesium chloride, 1 mM DTT, 20  $\mu\text{g}$  BSA, 10% glycerol, 0.5% Triton X-100) and poly dI-dC (1  $\mu\text{g}$ ) in a final volume of 20  $\mu\text{l}$ . The mixture was incubated at room temperature for 30 min. The gel was run as above, dried and exposed to film. For quantification, gels were analyzed by

phosphorimaging on a Molecular Dynamics Phosphor Imager (Amersham Biosciences). The identity of the NF- $\kappa$ B subunits was determined by including antibodies ([Santa Cruz Biotechnology, Santa Cruz, CA](#)) specific for p50 (sc-114), p52 (sc-848), Rel A (sc-372) or c-Rel (sc-71).

### Statistics

Statistics were calculated using Prism version 3.0 for Macintosh (GraphPad). Data are presented as means  $\pm$  SE. Data were analyzed using the Student's T-test or single factor ANOVA's with the Dunnet's multiple comparisons test.

## Results

### *LPS up-regulates cytokine gene induction in bone marrow stromal cells*

Bone marrow stromal cells support the growth and differentiation of early B cells (8). In previous studies, we demonstrated that cloned bone marrow stromal cell lines, including BMS2 cells, support the growth of primary [pre-B](#) and cloned ~~pre~~ and pro/pre-B cells and that treatment of the bone marrow stromal cells with AhR agonists alters their function (28-30, 32-34, 51). As a first step in determining if AhR ligands affect stromal cell cytokine production, baseline levels of cytokine mRNAs in BMS2 cells were determined by RNase protection assays.

RPAs with RNA from BMS2 cells consistently demonstrated significant expression of *IL-11*, ~~*IL-7*~~, *monocyte colony stimulating factor (M-CSF)*, *leukemia inhibitory factor (LIF)*, *IL-6*, *transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1)*, *TGF- $\beta$ 3* and *macrophage migration inhibitory factor (MIF)* mRNAs (Figure 1). Low but detectable levels of *IL-7* mRNA also were noted. Addition of the TLR-4 activator LPS consistently induced *GM-CSF*, *G-CSF*, *LIF* and *IL-6* mRNAs within 7 hours (Figure 1 and Table 1). As suggested by observations made with other stromal cell lines (44, 45), these studies demonstrate a potent innate inflammatory cytokine response against bacterial LPS.

### ***The effect of AhR agonists on LPS-mediated IL-6 mRNA induction***

To determine the effect of AhR ligands on LPS-induced cytokine induction, BMS2 cells were treated with vehicle (0.01% ethanol), 1  $\mu$ M DMBA, or 1 nM TCDD ~~1~~one hour prior to addition of 1-0  $\mu$ g/ml LPS. Cells were harvested 8 hours later, RNA was extracted, and assayed for cytokine mRNA levels by RPA.

Neither DMBA nor TCDD had a consistent effect on baseline levels of any cytokine mRNA assayed including IL-6 (Figure 2A and 2B). This result contrasts with that obtained with vascular endothelial cells in which AhR agonists (coplanar PCBs) alone induced IL-6 (52). As in previous experiments, LPS ~~induced~~ significantly induced *LIF*, and *IL-6* ~~(Figure 2A)~~ mRNAs (Figure 2A). (RNA encoding GM-CSF and G-CSF also were increased, although bands representing these RNAs were sometimes difficult to see when under-exposing blots to emphasize *IL-6* levels (e.g. Figure 2)).

Importantly, in five experiments, both DMBA and TCDD significantly reduced the LPS-mediated *IL-6* mRNA increase in BMS2 cells (Figure 2A and 2B;  $p < 0.05$ ). Other genes activated by LPS, i.e. *LIF*, *GM-CSF*, and *G-CSF*, were not affected by either of these AhR agonists (Figure 2A and data not shown) demonstrating the selectivity of AhR agonist-mediated effects on stromal cells.

Kinetics studies were performed using real-time PCR as a readout to quantify this change in *IL-6* mRNA levels. As seen in Figure 3, *IL-6* mRNA levels increased within 1 hr of LPS stimulation and peaked at the four hour time point. Inclusion of DMBA in the cultures reduced *IL-6* mRNA induction at every time point, including at 1 hour. These results are consistent with those obtained by RPA and demonstrate that AhR ligands suppress *IL-6* mRNA levels by approximately 40-60%.

### ***DMBA and TCDD significantly abrogate LPS-induced IL-6 production***

The results described above indicate consistent suppression of LPS-mediated *IL-6* mRNA induction. To determine if this decrease in steady state mRNA is reflected in a proportional decrease in secreted IL-6 protein levels, BMS2 cells were cultured for 18 hrs prior to addition of vehicle, 1  $\mu$ M

DMBA, or 1 nM TCDD. One hour later, cultures were treated with PBS (to assay effects of AhR ligands on background IL-6 levels) or challenged with 1.0 µg/ml LPS (i.e. time zero). Culture supernatants were harvested at time zero and 6-72 hours thereafter and assayed by ELISA for IL-6 levels.

Standardization with recombinant IL-6 indicated that this ELISA was able to detect as little as 15 pg/ml IL-6 (data not shown). Supernatants from BMS2 cultures established 18-24 hrs previously contained approximately 700 pg/ml IL-6 (Figure 4A). Addition of DMBA had no effect on IL-6 levels at any time point assayed. However, TCDD significantly ( $p < 0.05$ ) reduced the background level of IL-6 as early as 24 hrs after its addition to the cultures. Control experiments in which recombinant IL-6 was assayed in the presence of TCDD indicated that TCDD did not interfere with the IL-6-specific ELISA (data not shown). Therefore, TCDD is capable of reducing baseline IL-6 protein levels even though a reduction in baseline levels of *IL-6* mRNA was not evident (e.g. Figure 2B).

A significant increase in secreted IL-6 levels was seen as early as 6 hours after LPS treatment (Figure 4B). IL-6 levels continued to rise throughout the 72 hour period after LPS exposure. However, this LPS-dependent increase in secreted IL-6 was suppressed in cultures containing either DMBA or TCDD, with TCDD suppressing IL-6 levels as much as 88% at the 72 hour time point. Neither TCDD nor DMBA affected BMS2 proliferation as all cultures reached confluency at the same time (approximately 96 hrs after plating). These data demonstrate that activation of the AhR has a significant effect on the ability of bone marrow stromal cells to produce IL-6. Furthermore, the profound decrease in IL-6 protein levels in TCDD-exposed cells exhibiting only a 40-60% decrease in *IL-6* mRNA levels, suggests that at least this AhR ligand may suppress IL-6 production at more than just the RNA level.

***NF-κB activation correlates with IL-6 mRNA levels***

LPS-induced *IL-6* gene transcription in murine monocytes is controlled primarily by NF- $\kappa$ B, although other transcription factors may play minor roles (48). Since we and others have demonstrated interactions between the AhR and the NF- $\kappa$ B signaling pathways (46, 47, 52, 53), it was important to determine if activation of the AhR influenced NF- $\kappa$ B activity in bone marrow stromal cells. To this end, BMS2 cells were treated with vehicle or DMBA one hour prior to exposure to LPS as in previous experiments. Cells were harvested 16-24 hours later, nuclear proteins were extracted, and assayed in electromobility shift assays (EMSAs) for binding of nuclear proteins to a radiolabelled NF- $\kappa$ B probes consisting of the NF- $\kappa$ B-binding upstream regulatory element (URE) of the *c-myc* promoter (29).

Data presented in Figure 5A extend previous studies performed with monocytes (REF) by demonstrating that LPS induces NF- $\kappa$ B-DNA binding in bone marrow stromal cells. The ability to supershift NF- $\kappa$ B-DNA complexes in these EMSAs with antibodies specific for the p50 or RelA/p65 subunits of NF- $\kappa$ B (Figure 5B) indicates that LPS primarily activates a conventional NF- $\kappa$ B complex probably consisting of p50/RelA. While DMBA alone had no effect on the low baseline levels of NF- $\kappa$ B-DNA binding, it modestly suppressed the induction of NF- $\kappa$ B activity induced with LPS (Figure 5A). Quantification of normalized band densities indicated that DMBA significantly decreased LPS-induced NF- $\kappa$ B-DNA binding by 33% ( $p < 0.05$ ) (Figure 5C). These data demonstrate a correlation between NF- $\kappa$ B activity and *IL-6* mRNA levels.

## Discussion

The present studies were designed to evaluate the ability of AhR ligands to influence the elaboration of a spectrum of bone marrow stromal cell cytokines. In this system, exposing AhR<sup>+</sup> stromal cells to LPS mimicked the innate cytokine response to a gram-negative bacterial challenge. A large body of information now indicates that LPS exerts its inflammatory effects primarily through activation of Toll-like receptor-4 (TLR-4), a member of a family of pattern recognition receptors

expressed in virtually all tissue types (54, 55), including bone marrow stromal cells (56). TLRs play a critical role in innate immunity by responding directly to pathogen-associated molecules (57). Any environmental chemical capable of compromising this response has the potential to disrupt immunity to bacteria in specific and [the](#) regulation of inflammatory cytokines in general. The latter point is particularly important in the case of bone marrow stromal cells which use several cytokines, including IL-6, to regulate blood cell development.

Previously, we demonstrated that DMBA and TCDD, two prototypic AhR ligands, activate the AhR in BMS2 cells (28, 34). In studies presented here we demonstrate that both TCDD and DMBA suppress IL-6 production at least through the reduction of steady state *IL-6* mRNA levels. The specificity of this toxicity is underscored by the failure of these AhR agonists to affect either the baseline or LPS-induced levels of mRNAs specific for several other cytokines. This result also indicates that the observed changes in IL-6 do not result from overt toxicity or suppression of cell growth by AhR agonists.

Interestingly, TCDD reduced secretion of IL-6 to a greater extent (up to 88%) than might be expected from the RNA studies in which *IL-6* mRNA was only inhibited approximately 50%. This result could reflect the ability of TCDD to suppress IL-6 protein ~~transcription~~, secretion, or stability [in addition to lowering \*IL-6\* mRNA levels](#). Experiments are underway to evaluate these possibilities.

In our continuing investigations into interactions between AhR and NF- $\kappa$ B signaling pathways (47), we sought to determine if AhR signaling influences NF- $\kappa$ B activity in bone marrow stromal cells. An added incentive for these studies was provided by the observation that the *IL-6* gene promoter contains an NF- $\kappa$ B binding site which plays a major role in regulating LPS-induced *IL-6* transcription (58-60). Indeed, EMSAs indicated a profound increase in NF- $\kappa$ B activity following LPS exposure. This induction was suppressed modestly (33%) but significantly ( $p < 0.05$ ) by DMBA. These results suggest

a possible mechanism by which the AhR may regulate *IL-6* transcription, i.e. suppression of NF- $\kappa$ B signaling.

The AhR ligand-mediated decrease in NF- $\kappa$ B activity documented here differs from results obtained with other cell types. For example, coplanar PCBs, which can activate the AhR, increase IL-6 production in endothelial cells (61). Furthermore, TCDD increases both AP-1 and NF- $\kappa$ B binding to their cognate response elements in hepatoma cells (62, 63). In the latter example, NF- $\kappa$ B up-regulation involved an increase in binding of p50 homodimers to a consensus NF- $\kappa$ B nucleotide sequence. Unlike the p50/RelA-containing complexes implicated in the present studies (Figure 5B), p50 homodimers may actually block p50/p65-dependent gene activation (64). Indeed, p50/RelA dimers are well known to activate gene transcription (29, 50, 65, 66). Therefore, its suppression by AhR ligands is likely to have important biological consequences.

## **Conclusion**

The studies presented here demonstrate that prototypic AhR agonists, including an environmentally ubiquitous dioxin, specifically suppress the LPS-induced production of IL-6 by bone marrow stromal cells. The more potent AhR agonist, TCDD, also reduced baseline levels of IL-6 protein. Inhibition of IL-6 production in response to LPS occurs at least at the level of mRNA expression and may involve down-regulation of NF- $\kappa$ B, a transcription factor known to play a major role in regulating *IL-6* gene transcription. Collectively, the results demonstrate the potential for common environmental AhR ligands to compromise the ability of bone marrow stromal cells to respond to bacterial challenge and to support production of blood cell lineages.

**Table 1: LPS induces GM-CSF, G-CSF, LIF, and IL-6 mRNAs in bone marrow stromal cells**

Gene	N	Mean Fold Change <sup>a</sup> ± SE	P-value <sup>b</sup>
IL-11	4	1.3 ± 0.2	0.27
IL-7	5	0.9 ± 0.2	0.58
GM-CSF	3	Increased <sup>c</sup>	-
G-CSF	4	Increased <sup>c</sup>	-
LIF	5	5.7 ± 1.2 <sup>e</sup>	0.02*
IL-6	7	7.0 ± 1.5 <sup>e</sup>	0.01*
SCF	4	1.1 ± 0.1	0.40
M-CSF	5	1.0 ± 0.1	0.79
TGF-β1	2	1.3 ± 0.1	0.26
TGF-β3	2	0.9 ± 0.0	0.18
MIF	2	0.9 ± 0.1	0.52

<sup>a</sup> From the Ribonuclease Protection Assays (RPAs), mean fold-change is determined by normalizing the optical density band of the LPS-treated sample to the housekeeping gene GAPDH or L-32, then expressing it relative to the optical density of the normalized band from the vehicle-treated sample.

<sup>b</sup> Based on one-sample t-test to determine whether mean was significantly different from maximum LPS induction ( $\mu = 1.0$ ).

<sup>c</sup> Expression of GM-CSF and G-CSF was sometimes undetectable in unstimulated cells, thus the LPS-induced expression could not be described appropriately as "fold change"

\*  $p < 0.05$

## Figure Legends

**Figure 1: LPS induces GM-CSF, G-CSF, LIF, and IL-6 mRNAs in bone marrow stromal cells.**

BMS2 (bone marrow stromal) cells were cultured in the presence of 1 µg/ml LPS for eight hours. Cells were harvested, RNA was extracted, and mRNAs specific for several cytokines were assayed by RNase protection assays (RPAs). Representative data from two experiments (seven total) are presented.

**Figure 2. LPS-induced IL-6 mRNA induction is inhibited by DMBA and TCDD.** BMS2 cells were pre-treated with vehicle (0.01% ethanol), 1 µM DMBA, or 1 nM TCDD for one hour prior to challenge with 1 µg/ml LPS. Eight hours later, cells were harvested, RNA was extracted, and cytokine-specific mRNAs were assayed by RPA. (A) Representative data from two experiments (seven total) are presented. (B) Optical densities of bands corresponding to each cytokine mRNA were normalized to L-32 band densities and then expressed relative to the maximal response seen with LPS-plus-vehicle-treated cultures. Data represent the means ± SE from 4-5 independent experiments. One-sample t-tests were conducted to compare relative expressions to the theoretical  $\mu=1$  (maximum response in LPS-plus-vehicle-treated cultures). \* Significantly different from cultures treated with LPS and vehicle,  $p<0.05$ ; \*\* Significantly different from cultures treated with LPS and vehicle  $p<0.0001$ .

**Figure 3. Kinetics of DMBA-mediated suppression of IL-6 mRNA levels measured by-Real-time PCR.** BMS2 cells were pre-treated in triplicate with vehicle (0.01% ethanol) or 1 µM DMBA 1 hour prior to challenge with 1 µg/ml LPS. At various points thereafter, cells in individual wells were harvested separately, was-RNA was extracted, and cytokine-specific mRNAs were quantified by real-time PCR. Data represent means ± SE of IL-6 mRNA levels (number of PCR cycles for half- maximal

amplification) normalized to  $\beta$ -actin levels according to the Comparative C<sub>T</sub> Method (PE Applied Biosystems) and expressed relative to maximal *IL-6* levels in LPS-plus-vehicle-treated cultures at the 4 hour time point. Representative data from a total of three experiments are shown.

**Figure 4. DMBA and TCDD suppress LPS-induced *IL-6* protein production.** BMS2 cells were cultured for 18-24 hours prior to treatment with vehicle (0.01% ethanol), 1  $\mu$ M DMBA, or 1 nM TCDD. One hour later, cultures were challenged with 1  $\mu$ g/ml LPS as indicated. Supernatants were sampled at various times thereafter and assayed by ELISA for *IL-6* protein levels. *IL-6* concentrations were determined by comparison to a recombinant *IL-6* standard curve using recombinant *IL-6*. (A) *IL-6* levels in groups not challenged with LPS. \*Significantly different from vehicle control,  $p < 0.05$  (one-way ANOVA and Dunnett's multiple comparisons test). \*\*Significantly different from vehicle control,  $p < 0.01$ . (B) *IL-6* levels in groups challenged with LPS. \*Significantly different from corresponding LPS-plus-vehicle wells ( $p < 0.05$ ). \*\*Significantly different from vehicle or vehicle-plus-LPS wells,  $p < 0.01$ .

**Figure 5: DMBA inhibits LPS-induced NF- $\kappa$ B-DNA binding.** BMS2 cells were left untreated (naïve) or were pre-treated with vehicle (0.01% ethanol) or 1  $\mu$ M DMBA for one hour prior to challenge with 1  $\mu$ g/ml LPS. Cells were harvested 16 and 24 hrs later; Nuclear proteins were extracted, and proteins analyzed by EMSA for binding to an NF- $\kappa$ B probe derived from the *c-myc* upstream regulatory element (URE). (A) Data from a representative experiment (three total) are presented. An arrow indicates the NF- $\kappa$ B-specific band. (B) EMSAs (Jennifer – which timepoint?) were performed with nuclear extracts from BMS2 cells treated for 30 min?? (I'll check in the am) hrs with LPS in the presence of antibodies specific for the p50, p53, RelA (p65), or c-Rel subunits of NF- $\kappa$ B.

(C) Quantification of NF- $\kappa$ B-DNA binding in nuclear extracts treated for 16-24 hrs. (~~Jennifer is this the pooled data for 16 and 24 hrs?~~) — NF- $\kappa$ B band densities were normalized to untreated controls within the same experiments. \*\*Significantly different from LPS- plus vehicle-treated control, p<0.05 ~~treated vehicle control, p<0.05 (one-way ANOVA and Dunnett's multiple comparisons test)~~ (pPaired T-test, Vh+LPS vs. DMBA+LPS).

## Abbreviations

AhR, aryl hydrocarbon receptor/transcription factor; B[a]P, benzo[a]pyrene; DMBA, 7,12-dimethylbenz[a]anthracene; G-CSF, granulocyte colony stimulating factor; GM-CSF, granulocyte/monocyte colony stimulating factor; HAH, halogenated aromatic hydrocarbons; IFN- $\gamma$ , interferon- $\gamma$ ; LIF, leukemia inhibitory factor; LPS, Staphylococcus-derived lipopolysaccharide; MIF, macrophage migration inhibitory factor; PAH, polycyclic aromatic hydrocarbon(s); TCDD-2,3,7,8-tetrachlorodibenzo-p-dioxin; TLR, Toll like receptor(s); TNF, tumor necrosis factor; TGF, transforming growth factor

## Competing Interests

Supported by NIH RO1-ES06086, Superfund Basic Research Grant #1P42ES-07381, and PO1 HL 68705

## Authors' Contributions

B.A.J. designed and performed experiments summarized in Figures 1-4 and Table 1 with the assistance of R.J.L. and shared responsibilities for planning the project and drafting this manuscript with D.H.S.. J.J.S. performed experiments summarized in Figure 5. D.H.S. proposed the general field of study, participated in the design of experiments, supervised data interpretation, and obtained funding to support the work.

**Acknowledgements**

The authors thank Ilga Wohlrab for performing all the administrative duties related to this project and to Maryam Shansab for assisting in manuscript preparation.

## REFERENCES

1. Kittler, E. L., H. McGrath, D. Temeles, R. B. Crittenden, V. K. Kister, and P. Quesenberry. 1992. Biologic significance of constitutive and sublethal growth factor production by bone marrow stroma. *Blood* 79:3168.
2. Robledo, M. M., and J. Teixido. 1997. TGF- $\beta$ 1 binding proteins on human bone marrow stromal cells. *Leuk Lymphoma* 27:509.
3. Gupta, D., S. P. Treon, Y. Shima, T. Hideshima, K. Podar, Y. T. Tai, B. Lin, S. Lentzsch, F. E. Davies, D. Chauhan, R. L. Schlossman, P. Richardson, P. Ralph, L. Wu, F. Payvandi, G. Muller, D. I. Stirling, and K. C. Anderson. 2001. Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications. *Leukemia* 15:1950.
4. Akiyama, M., T. Hideshima, T. Hayashi, Y. T. Tai, C. S. Mitsiades, N. Mitsiades, D. Chauhan, P. Richardson, N. C. Munshi, and K. C. Anderson. 2002. Cytokines modulate telomerase activity in a human multiple myeloma cell line. *Cancer Res* 62:3876.
5. Petrucci, M., M. R. Ricciardi, C. Gregorj, L. Ciapponi, R. Savino, G. Ciliberto, and A. Tafuri. 2002. Effects of IL-6 variants in multiple myeloma: growth inhibition and induction of apoptosis in primary cells. *Leuk Lymphoma* 43:2369.
6. Aoyama, K., K. Oritani, T. Yokota, J. Ishikawa, T. Nishiura, K. Miyake, Y. Kanakura, Y. Tomiyama, P. W. Kincade, and Y. Matsuzawa. 1999. Stromal cell CD9 regulates differentiation of hematopoietic stem/progenitor cells. *Blood* 93:2586.
7. Hirose, J., T. Kouro, H. Igarashi, T. Yokota, N. Sakaguchi, and P. W. Kincade. 2002. A developing picture of lymphopoiesis in bone marrow. *Immunol Rev* 189:28.
8. Kincade, P. W., G. Lee, C. E. Pietrangeli, S.-I. Hayashi, and J. M. Gimble. 1989. Cells and molecules that regulate B lymphopoiesis in bone marrow. *Ann. Rev. Immunol.* 7:111.

9. Pietrangeli, C. E., S. Hayashi, and P. W. Kincade. 1988. Stromal cell lines which support lymphocyte growth: characterization, sensitivity to radiation and responsiveness to growth factors. *Eur J Immunol* 18:863.
10. Billiau, A. 1986. BSF-2 is not just a differentiation factor. *Nature* 324:415.
11. Gimble, J. M., C. Pietrangeli, A. Henley, M. A. Dorheim, J. Silver, A. Namen, M. Takeichi, C. Goridis, and P. W. Kincade. 1989. Characterization of murine bone marrow and spleen-derived stromal cells: analysis of leukocyte marker and growth factor mRNA transcript levels. *Blood* 74:303.
12. Taga, T., and T. Kishimoto. 1992. Role of a two-chain IL-6 receptor system in immune and hematopoietic cell regulation. *Crit Rev Immunol* 11:265.
13. Kopf, M., A. Ramsay, F. Brombacher, H. Baumann, G. Freer, C. Galanos, J. C. Guteirrez-Ramos, and G. Kohler. 1995. Pleiotropic defects of IL-6-deficient mice including early hematopoiesis, T and B cell function, and acute phase responses. *Ann N Y Acad Sci* 762:308.
14. Kawano, M. M., H. Ishikawa, N. Tsuyama, S. Abroun, S. Liu, F. J. Li, K. Otsuyama, and X. Zheng. 2002. Growth mechanism of human myeloma cells by interleukin-6. *Int J. Hematol* 76:329.
15. Wols, H. A. M., G. H. Underhill, G. S. Kansas, and P. L. Witte. 2002. The role of bone marrow-derived stromal cell in the maintenance of plasma cell longevity. *J. Immunol* 169:4213.
16. Tu, C. X., J. S. Gu, and X. R. Lin. 2003. Increased interleukin-6 and granulocyte-macrophage colony stimulating factor levels in the sera of patients with non-segmental vitiligo. *J Dermatol Sci* 31:73.
17. Finck, B. K., B. Chan, and D. Wofsy. 1994. Interleukin 6 promotes murine lupus in NZB/NZW F1 mice. *J Clin Invest* 94:585.

18. McMurray, R. W., R. W. Hoffman, W. Nelson, and S. E. Walker. 1997. Cytokine mRNA expression in the B/W mouse model of systemic lupus erythematosus--analyses of strain, gender, and age effects. *Clin Immunol Immunopathol* 84:260.
19. Nakahara, H., J. Song, M. Sugimoto, K. Hagihara, T. Kishimoto, K. Yoshizaki, and N. Nishimoto. 2003. Anti-interleukin-6 receptor antibody therapy reduces vascular endothelial growth factor production in rheumatoid arthritis. *Arthritis Rheum* 48:1521.
20. Heinrich, P. C., I. Behrmann, S. Haan, H. M. Hermanns, G. Muller-Newen, and F. Schaper. 2003. Principles of interleukin (IL)-6-type cytokine signalling and its regulation. *Biochem J* 374:1.
21. Jourdan, M., J. L. Veyrune, J. D. Vos, N. Redal, G. Couderc, and B. Klein. 2003. A major role for Mcl-1 antiapoptotic protein in the IL-6-induced survival of human myeloma cells. *Oncogene* 22:2950.
22. Hirata, T., C. Shimazaki, T. Sumikuma, E. Ashihara, H. Goto, T. Inaba, Y. Koishihara, and M. Nakagawa. 2003. Humanized anti-interleukin-6 receptor monoclonal antibody induced apoptosis of fresh and cloned human myeloma cells in vitro. *Leuk Res* 27:343.
23. Jee, S. H., S. C. Shen, H. C. Chiu, W. L. Tsai, and M. L. Kuo. 2001. Overexpression of interleukin-6 in human basal cell carcinoma cell lines increases anti-apoptotic activity and tumorigenic potency. *Oncogene* 20:198.
24. An, J., Y. Sun, R. Sun, and M. B. Rettig. 2003. Kaposi's sarcoma-associated herpesvirus encoded vFLIP induces cellular IL-6 expression: the role of the NF-kappaB and JNK/AP1 pathways. *Oncogene* 22:3371.
25. Kallen, K. J. 2002. The role of transsignalling via the agonistic soluble IL-6 receptor in human diseases. *Biochim Biophys Acta* 1592:323.

26. Heidel, S. M., K. Holston, J. T. Buters, F. J. Gonzalez, C. R. Jefcoate, and C. J. Czuprynski. 1999. Bone marrow stromal cell cytochrome P4501B1 is required for pre-B cell apoptosis induced by 7,12-dimethylbenz[a]anthracene. *Mol Pharmacol* 56:1317.
27. Page, T. J., S. O'Brien, C. R. Jefcoate, and C. J. Czuprynski. 2002. 7,12-Dimethylbenz[a]anthracene induces apoptosis in murine pre-B cells through a caspase-8-dependent pathway. *Mol Pharmacol* 62:313.
28. Mann, K. K., R. A. Matulka, B. P. Lawrence, N. I. Kerkvliet, and D. H. Sherr. 1999. The role of cytochrome P-450 enzymes in 7,12-dimethylbenz[a]anthracene-induced apoptosis. *Toxicol. Appl. Pharmacol.* 161:10.
29. Mann, K. K., S. Doerre, D. H. Sherr, and S. Quadri. 2001. The Role of NF- $\kappa$ B as a survival factor in environmental chemical-induced pre-B cell apoptosis. *Molec. Pharmacol.* 59:302.
30. Near, R. I., K. K. Mann, R. A. Matulka, A. M. Shneider, S. U. Gogate, A. F. Trombino, and D. H. Sherr. 1999. Regulation of pre-B cell apoptosis by aryl hydrocarbon receptor/transcription factor-expressing stromal/adherent cells. *Proc. Soc. Exp. Biol. Med.* 221:242.
31. Allan, L. L., K. K. Mann, J. L. Schlezinger, and D. H. Sherr. 2003. Bone marrow interactions in polycyclic aromatic hydrocarbon-induced pro/pre-B cell apoptosis. *Toxicol. Sci. In Press.*
32. Ryu, H.-Y., K. K. Mann, J. J. Schlezinger, B. Jensen, and D. H. Sherr. 2003. Environmental chemical-induced pro/pre-B cell apoptosis: Analysis of c-Myc, p27<sup>Kip1</sup>, and p21<sup>WAF1</sup> reveals a death pathway distinct from clonal deletion. *J. Immunol.* 170:4897.
33. Yamaguchi, K., R. A. Matulka, A. Shneider, P. Toselli, A. F. Trombino, S. Yang, L. J. Hafer, K. K. Mann, X.-J. Tao, J. L. Tilly, R. Near, and D. H. Sherr. 1997. Induction of pre-B cell apoptosis by 7,12 dimethylbenz[a]anthracene in long term bone marrow cultures. *Toxicol. Appl. Pharmacol.* 147:190.

34. Yamaguchi, K., R. I. Near, R. A. Matulka, A. Shneider, P. Toselli, A. F. Trombino, and D. H. Sherr. 1997. Activation of the aryl hydrocarbon receptor/transcription factor and stromal cell-dependent pre-B cell apoptosis. *J. Immunol.* 158:2165.
35. Denison, M., L. Vella, and A. Okey. 1986. Structure and function of the Ah Receptor for 2,3,7,8-tetrachlorodibenzo-p-dioxin. *The Journal of Biological Chemistry* 261:3987.
36. Hogenesch, J. B., W. K. Chan, V. H. Jackiw, R. C. Brown, Y. Z. Gu, M. Pray-Grant, G. H. Perdew, and C. A. Bradfield. 1997. Characterization of a subset of the basic-helix-loop-helix-PAS superfamily that interacts with components of the dioxin signaling pathway. *J Biol Chem* 272:8581.
37. Perdew, G. 1988. Association of the Ah receptor with the 90-kDa heat shock protein. *The Journal of Biological Chemistry* 263:13802.
38. Murante, F. G., and T. A. Gasiewicz. 2000. Hemopoietic progenitor cells are sensitive targets of 2,3,7,8-tetrachlorodibenzo-p-dioxin in C57BL/6J mice. *Toxicol Sci* 54:374.
39. Staples, J. E., F. G. Murante, N. C. Fiore, T. A. Gasiewicz, and A. E. Silverstone. 1998. Thymic alterations induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin are strictly dependent on aryl hydrocarbon receptor activation in hemopoietic cells. *J Immunol* 160:3844.
40. Schwartz, G. G. 1997. Multiple myeloma: clusters, clues, and dioxins. *Canc. Epi., Biomarkers, Prev.* 6:49.
41. Bertazzi, P. A., A. C. Pesatori, D. Consonni, A. Tironi, M. T. Landi, and C. Zocchetti. 1993. Cancer incidence in a population accidentally exposed to 2,3,7,8-tetrachlorodibenzo-para-dioxin. *Epidemiol.* 4:398.
42. Bertazzi, P. A., C. Zocchetti, S. Guercilena, D. Consonni, A. Tironi, M. T. Landi, and A. C. Pesatori. 1997. Dioxin exposure and cancer risk: A 15-year mortality study after the "Seveso accident". *Epidemiol.* 8:646.

43. Schilling, J. D., S. M. Martin, C. S. Hung, R. G. Lorenz, and S. J. Hultgren. 2003. Toll-like receptor 4 on stromal and hematopoietic cells mediates innate resistance to uropathogenic *Escherichia coli*. *Proc Natl Acad Sci U S A* 100:4203.
44. Lorgeot, V., F. Rougier, P. Fixe, E. Cornu, V. Praloran, and Y. Denizot. 1997. Spontaneous and inducible production of leukaemia inhibitory factor by human bone marrow stromal cells. *Cytokine* 9:754.
45. Rougier, F., E. Cornu, V. Praloran, and Y. Denizot. 1998. Il-6 and IL-8 production by human bone marrow stromal cells. *Cytokine* 10:93.
46. Tian, Y., S. Ke, M. S. Denison, A. B. Rabson, and M. A. Gallo. 1999. Ah receptor and NF- $\kappa$ B interactions, a potential mechanism for dioxin toxicity. *J. Biol. Chem.* 274:510.
47. Kim, D. W., L. Gazourian, S. A. Quadri, D. H. Sherr\*, and G. E. Sonenshein\*. 2000. The aryl hydrocarbon receptor/transcription factor (AhR) and the Rel A nuclear factor- $\kappa$ B subunit cooperate to transactivate the *c-myc* promoter. *Oncogene* 19:5498.
48. Dendorfer, U., P. Oettgen, and T. A. Libermann. 1995. Interleukin-6 Gene Expression by Prostaglandins and Cyclic AMP Mediated by Multiple Regulatory Elements. *Am J Ther* 2:660.
49. Schlezinger, J. J., B. A. Jensen, K. K. Mann, H. Y. Ryu, and D. H. Sherr. 2002. Peroxisome Proliferator-Activated Receptor  $\gamma$ -Mediated NF- $\kappa$ B Activation and Apoptosis in Pre-B Cells. *J Immunol* 169:6831.
50. Duyao, M., A. Buckler, and G. Sonenshein. 1990. Interaction of an NF- $\kappa$ B-like factor with a site upstream of the *c-myc* promoter. *Proc. Natl. Acad. Sci. USA* 87:4727.
51. Quadri, S., A. Qadri, K. L. Mann, and D. H. Sherr. 2000. The bioflavonoid galangin blocks aryl hydrocarbon receptor (AhR) activation and polycyclic aromatic hydrocarbon-induced pre-B cell apoptosis. *Molec. Pharmacol.* 58:515.

52. Hennig, B., P. Meerarani, R. Slim, M. Toborek, A. Daugherty, A. E. Silverstone, and L. W. Robertson. 2002. Proinflammatory properties of coplanar PCBs: in vitro and in vivo evidence. *Toxicol Appl Pharmacol* 181:174.
53. Yao, Y., A. Hoffer, C. Y. Chang, and A. Puga. 1995. Dioxin activates HIV-1 gene expression by an oxidative stress pathway requiring a functional cytochrome P450 CYP1A1 enzyme. *Environ Health Perspect* 103:366.
54. Chaudhary, P. M., C. Ferguson, V. Nguyen, O. Nguyen, H. F. Massa, M. Eby, A. Jasmin, B. J. Trask, L. Hood, and P. S. Nelson. 1998. Cloning and characterization of two Toll/Interleukin-1 receptor-like genes TIL3 and TIL4: evidence for a multi-gene receptor family in humans. *Blood* 91:4020.
55. Rock, F. L., G. Hardiman, J. C. Timans, R. A. Kastelein, and J. F. Bazan. 1998. A family of human receptors structurally related to *Drosophila* Toll. *Proc Natl Acad Sci USA* 95:588.
56. Kikuchi, T., T. Matsuguchi, N. Tsuboi, A. Mitani, S. Tanaka, M. Matsuoka, G. Yamamoto, T. Hihikawa, T. Noguchi, and Y. Yoshikai. 2001. Gene expression of osteoclast differentiation factor is induced by lipopolysaccharide in mouse osteoblasts via Toll-like receptors. *J Immunol* 166:3574.
57. Zhang, G., and S. Ghosh. 2001. Toll-like receptor-mediated NF-kappaB activation: a phylogenetically conserved paradigm in innate immunity. *J Clin Invest* 107:13.
58. Dendorfer, U., P. Oettgen, and T. A. Libermann. 1994. Multiple regulatory elements in the interleukin-6 gene mediate induction by prostaglandins, cyclic AMP, and lipopolysaccharide. *Mol Cell Biol* 14:4443.
59. Baccam, M., S. Y. Woo, C. Vinson, and G. A. Bishop. 2003. CD40-mediated transcriptional regulation of the IL-6 gene in B lymphocytes: involvement of NF-kappa B, AP-1, and C/EBP. *J Immunol* 170:3099.

60. Zerbini, L. F., Y. Wang, J. Y. Cho, and T. A. Libermann. 2003. Constitutive activation of nuclear factor kappaB p50/p65 and Fra-1 and JunD is essential for deregulated interleukin 6 expression in prostate cancer. *Cancer Res* 63:2206.
61. Henning, B., P. Meerarani, R. Slim, M. Toberek, A. Daugherty, A. E. Silverstone, and L. W. Robertson. 2002. Proinflammatory properties of coplanar PCBs: In vitro and in vivo evidence. *Toxicol Appl Pharmacol* 181:174.
62. Puga, A., D. Nebert, and F. Carrier. 1992. Dioxin induces expression of *c-fos* and *c-jun* proto-oncogenes and a large increase in transcription factor AP-1. *DNA and Cell Biol.* 11:269.
63. Puga, A., S. J. Barnes, T. P. Dalton, C.-y. Chang, E. S. Knudsen, and M. A. Maier. 2000. Aromatic hydrocarbon receptor interaction with the retinoblastoma protein potentiates repression of E2F-dependent transcription and cell cycle arrest. *J. Biol. Chem.* 275:2943.
64. Kang, S. M., A. C. Tran, M. Grilli, and M. J. Lenardo. 1992. NF-kappa B subunit regulation in nontransformed CD4+ T lymphocytes. *Science* 256:1452.
65. Duckett, C. S., N. D. Perkins, T. F. Kowalik, R. M. Schmid, E. S. Huang, J. A.S. Baldwin, and G. J. Nabel. 1993. Dimerization of NF- $\kappa$ B2 with RelA(p65) regulates DNA binding, transcriptional activation, and inhibition by an I $\kappa$ B- $\alpha$  (MAD-3). *Mole. Cell. Biol.* 13:1315.
66. Gilmore, T. D., M. Koedood, K. A. Piffat, and D. W. White. 1996. Rel/NF- $\kappa$ B/I $\kappa$ B proteins and cancer. *Oncogene* 13:1367.











