

Title: Respiratory symptoms in children living near busy roads and their relationship to car and truck traffic: results of an Italian multicenter study (SIDRIA 2)

Authors: G. Berti¹, E. Migliore², C. Galassi², N. Pearce³⁻⁴, F. Forastiere⁵, L. Armenio⁶, A. Biggeri⁷⁻⁸, L. Bisanti⁹, M. Bugiani¹⁰, E. Cadum¹, E. Chellini¹¹, V. Dell'Orco¹², G. Giannella¹³, P. Sestini¹⁴, G. Corbo¹⁵, R. Pistelli¹⁵, G. Viegi¹⁶, G. Ciccone² and SIDRIA-2 Collaborative Group¹⁷

Istitutional addresses

¹ Regional Environmental Protection Agency, Piedmont Region, Via Sabaudia 164, 10095 Grugliasco (Turin), Italy

² Cancer Epidemiology Unit, AOU San Giovanni Battista Hospital - Center for Cancer Prevention (CPO Piedmont) and University of Turin, Via Santena 7, 10126 Turin, Italy

³ Centre for Public Health Research, Massey University Wellington Campus, PO Box 756 Wellington 6140 NZ

⁴ Department of Biomedical Sciences and Human Oncology, University of Turin, Via Santena 7, 10126 Turin, Italy

⁵ Department of Epidemiology, Rome/E Local Health Authority, Via di S. Costanza 53, 00198 Rome, Italy

⁶ I Pediatric Clinic, University of Bari, Piazza G. Cesare 11, 70124 Bari, Italy

⁷ Department of Statistics, University of Florence, Viale Morgagni 59, 50134 Florence, Italy

⁸ Unit of Biostatistics, Istituto per lo Studio e la Prevenzione Oncologica (ISPO), Via San Salvi 12, 50135 Florence, Italy

⁹ Epidemiology Unit, Local Health Authority, Corso Italia 19, 20122 Milan, Italy

¹⁰ Unit of Pneumology and Allergology, Local Health Authority 2, Lungo Dora Savona 26, 10152 Turin, Italy

¹¹ Unit of Environmental and Occupational Epidemiology, Istituto per lo Studio e la Prevenzione Oncologica (ISPO), Via San Salvi 12, 50135 Florence, Italy

¹² Department of prevention, Rome/G Local Health Authority - Tivoli Corso Garibaldi 7, 00034 Colleferro (Rome), Italy

¹³ Unit of Preventive Medicine, Local Health Authority, Via Trento 6, 46100 Mantova, Italy

¹⁴ Institute of Respiratory Diseases, University of Siena, Viale Bracci 3, 53100 Siena, Italy

¹⁵ Department of Respiratory Physiology, Catholic University of Rome, Largo F.Vito 1, 00168 Rome, Italy

¹⁶ CNR Institutes of Biomedicine and Molecular Immunology, Palermo, and of Clinical Physiology, Pisa. Via Ugo La Malfa 153, 90146 Palermo, Italy

¹⁷ SIDRIA-2 Collaborative Group: G. Ciccone, E. Migliore, D. Mirabelli, G. Berti, E. Cadum, M. Bugiani, P. Piccioni (Turin); L. Bisanti, A. Russo, F. Rusconi, M. Bellasio, V. Gianelle (Milan); S. Piffer, L. Battisti, D. Kaisermann, M. Gentilini (Trento); G. Giannella, F. Talassi (Mantova); C. Galassi, N. Caranci, G. Frasca, M. Biocca, E. De Munari (Emilia-Romagna); E. Chellini, E. Lombardi, A. Biggeri, C. Gabellini, D. Grechi (Florence); M.G. Petronio (Empoli); P. Sestini (Siena); G. Viegi, M. Simoni (Pisa); F. Forastiere, M. De Sario, R. Pistelli, G. Corbo, E. Bonci, L. Indinnimeo, V. Dell'Orco, N. Agabiti (Rome); L. Armenio, L. Brunetti, M. Cavone, M. Lospalluti, M. Massagli, G. Polieri, D. Rizzi, F. Rana, M. Rana (Bari); S. La Grutta (Palermo)

E-mail addresses:

GB: g.ber ti@arpa.piemonte.it

EM: enrica.migliore@cpo.it

CG: claudia.galassi@cpo.it

NP: n.e.pearce@massey.ac.nz

FF: forastiere@asplazio.it

LA: l.armenio@pediatria3.uniba.it

AB: abiggeri@ds.unifi.it

LB: lbisanti@asl.milano.it

MB: m.bugiani@aslto4.it

EC: e.cadum@arpa.piemonte.it;

ECh: e.chellini1@cspo.it;

VDO: valerio.dellorco@tiscali.it;

GG: gabriele.giannella@aslmn.it

PS: sestini@unisi.it;

GC: gmcorbo@yahoo.com;

RP: pneumologia@h-columbus.it;

GV: viegig@ifc.cnr.it;

GC: gianni.ciccone@cpo.it

Corresponding author:

Enrica Migliore

Cancer Epidemiology Unit,

AOU San Giovanni Battista Hospital - Center for Cancer Prevention (CPO Piedmont) and

University of Turin

Via Santena 7

10126 Torino - ITALY

Tel. +39 011 6336856

Fax +39 011 6334664

e-mail enrica.migliore@cpo.it

Abstract:

Background

Epidemiological studies have provided evidence that exposure to vehicular traffic increases the prevalence of respiratory symptoms and may exacerbate pre-existing asthma in children

The aim of this study was to investigate whether there were specific effects of cars and trucks traffic on current asthma symptoms (i.e. wheezing) and cough or phlegm.

Methods

The survey was conducted in 2002 in 12 centers of Northern, Center and Southern Italy, different in size, climate, latitude and level of urbanization. Standardized questionnaires were used to collect information on health outcomes and exposure to traffic among 6-7 and 13-14 years old children and adolescents. A questionnaire filled in by parents was available for 33632 subjects (response rate: 86.7%).

Results

Overall traffic density was weakly associated with asthma symptoms but there was a stronger association with cough or phlegm (high traffic density OR=1.24; 95% CI: 1.04, 1.49). Car and truck traffic were independently associated with cough or phlegm. The strongest associations were found when asthmatic symptoms and cough or phlegm were combined as the final outcome (for high traffic density OR=1.52; 95% CI: 1.17, 1.96).

Conclusions

Children living near streets with intense traffic are at higher risk for respiratory effects. Traffic exposure is mainly associated with symptoms of cough or phlegm, with or without asthma symptoms.

Background

Vehicular traffic is a major source of outdoor air pollution. Several studies have reported associations between exposure to traffic pollutants in the zone of residence and increased frequency of respiratory tract illnesses [1-10]. A specific role of diesel exhaust from heavy traffic has been suggested in some of these studies [4, 5, 7], and airway inflammation due to exposure to diesel exhaust seems the likely biological mechanism [11, 12]. With respect to the type of respiratory disorder, consistent associations have been found between exposure to traffic fumes and bronchitis symptoms, while the role of these exposures in the etiology of asthma is still unclear [13].

The SIDRIA project - Studi Italiani sui Disturbi Respiratori nell'Infanzia e l'Ambiente [14-16] - is a large multi-center, population-based study conducted in the framework of the International Study of Asthma and Allergies in Childhood (ISAAC) [17]. The findings from SIDRIA-1, conducted in 1994-1995, showed a positive association between indicators of air pollution from heavy vehicular traffic in the street of residence and a wide range of respiratory disorders in children living in highly urbanized areas [18]. However, it was not possible to assess the independent effects of car and truck traffic in that study as the information was not available.

A second phase of SIDRIA was conducted in 2002 to evaluate time trends in the prevalence of respiratory disorders in childhood, according to the ISAAC Phase III protocol [19], to confirm the role of several potential risk factors identified in SIDRIA-1 and to explore some new hypotheses. In this paper, based on the data from SIDRIA-2, we present an analysis of the relation between indicators of road traffic pollution and several chronic respiratory symptoms. The main purpose of this analysis was to confirm previous results and to evaluate the possible independent respiratory effects of each type of traffic exposure (truck traffic or car traffic). Since chronic cough may be a manifestation of asthma or it may occur also in otherwise healthy children mainly as a result of "back to back" respiratory infections or of recurrent or chronic rhino sinusitis with postnasal drip

[20], we have tried to distinguish the risk factors of asthma symptoms and of symptoms of persistent cough or sputum production (cough or phlegm).

Methods

The SIDRIA-2 study design has been described elsewhere [15, 16, 21] and it will be only summarized here. The survey was conducted in 2002, between January and May, in 12 centers (Bari, Colleferro, Emilia-Romagna, Empoli, Florence, Mantova, Milan, Palermo, Rome, Siena, Turin, Trento) of Northern, Center and Southern Italy, different in size, climate, latitude and level of urbanization. Eight of these centers had already participated in SIDRIA-1 [18]. The protocol of the study was approved by Ethics Committee of the Catholic University in Rome.

The sample included 22442 children (6-7 years old) and 16336 adolescents (13-14 years old) attending respectively the first two grades of the primary school and the last year of the middle school. The primary sampling units were schools, both public and private, weighted for the number of attending subjects. Each center contributed with at least 1000 subjects for each age group. To collect information on the medical history of the children, we used standardized, self administered questionnaires that included also the relevant ISAAC-Phase III questions on asthma, rhinitis and eczema symptoms, and questions on various known or suspected risk factors. For children (6-7 years old), all questionnaires were completed by parents. According to the standard ISAAC protocol [17], the questionnaires for adolescents were filled by adolescents themselves. However, in the SIDRIA study, another questionnaire, including questions on both symptoms and risk factors, was also completed by the adolescents' parents. For reasons of consistency and comparability between age groups, the current analyses are based on the parental questionnaires for both age-groups.

Children with “asthma symptoms” were defined as those reporting in the past 12 months at least one of the following: one or more wheezing episodes, wheeze with exercise, morning chest tightness; or if they reported night dry cough in the last 12 months and had a reporting of lifetime asthma; or those reporting treatment for medically diagnosed asthma or had a hospital admission for asthma in the last 12 months; or if they reported a life time asthma and a positive answer to the question “Is your child still suffering from asthma?”.

Children were defined as having “severe asthma” if in the past 12 months at least one of the following were reported: 4 or more wheezing attacks, waking at night with wheezing one or more times a week, an attack severe enough to limit speech to only one or two words at a time between breaths, or a hospital admission for asthma.

Children were defined as having “cough or phlegm” if they reported cough or phlegm for at least 4 days a week (in the absence of a cold) for one or more months a year.

Questions on traffic included a parental subjective evaluation of traffic density in the zone of residence (“absent”, “low”, “moderate” or “high”) and of the daily frequency of passing cars and trucks in the street of residence (“never or seldom”, “sometimes”, “frequently”, “continuously”). The exact wording is reported in the Appendix 1 (Table 1).

Odds ratio (OR) and 95% confidence intervals (95% CI) were estimated with multiple logistic regression analyses. The basic analyses were for “asthma symptoms” and “cough or phlegm”, but we also conducted multinomial logistic regression analyses comparing: i) those with asthma symptoms without cough or phlegm; ii) those with cough or phlegm without asthma symptoms; and iii) those with the combination of the two conditions, with the remaining subjects, negative for both symptoms. For “cough or phlegm”, the analyses focused on the possible independent effect of cars and trucks transit in the street of residence. In some of the analyses, in order to obtain sufficient numbers in each category, car transit was recoded into three categories (“absent/sometimes”,

“frequently” and “continuously”) and truck transit was also recoded into three categories (“absent”, “sometimes” and “frequently/continuously”).

Potential confounding factors included in the multiple logistic regression models were: sex, age, parental asthma or allergy (rhinitis or eczema), parental education (higher educational level between parents as a proxy of socioeconomic status – SES), passive smoke at home (at least one smoker -mother, father or others- in the household), indoor mould/dampness, floor of the apartment, change of residence and study area. The effect of cluster sampling by school was considered and each regression model was adjusted for autocorrelation within schools. In order to explore the role of potential effect modifiers, we performed subgroup analyses for different factors (age, gender, latitude, parental education, smoking, parental asthma or allergies, level of urbanization, indoor mould/dampness, floor of the apartment, change of residence), in each case considering the associations of truck traffic exposure (frequent or continuous vs never) and the investigated respiratory symptoms. Statistical significance of the interaction terms were evaluated. All analyses were conducted using STATA 9 (Stata Corporation, College Station, Texas).

Results

The parental questionnaire was completed for 20016 6-7 years old (response rate 89.2%) and for 13616 13-14 years old (response rate 83.3%). Table 1 shows the combined prevalence of asthma symptoms and cough or phlegm. Overall, 13.5% of children (95% CI: 13.2%, 13.9%) were reported to have asthma symptoms, and 6.8% (95% CI: 6.6%, 7.1%) to have cough or phlegm. For 2.9% (95% CI: 2.7%, 3.0%) of the subjects, cough or phlegm and asthma symptoms were concomitant. The frequency of each traffic indicator is presented in Table 2 of the appendix.

The prevalences of respiratory symptoms in the various subgroups are shown in Table 2. Respiratory symptoms were more frequent if a parental history of asthma or allergies was reported,

among subjects exposed to passive tobacco smoke and in the presence of moulds/dampness in the child's bedroom. The prevalence of respiratory symptoms increased inversely with parental education, and directly with urbanization level.

Table 3 shows the associations of traffic indicators with asthma symptoms and with cough or phlegm. Reported high traffic density, continuous car transit and continuous truck transit in the street of residence were weakly associated with asthma symptoms. There were stronger associations of reported high traffic density, continuous car transit and continuous truck transit with cough or phlegm. The associations were generally stronger for truck transit than for car transit.

For comparison with the literature, we present the results of the associations for different asthma symptoms in Table 3 of the appendix. In the same table, we also report the associations separately for light and severe asthma symptoms, and for symptoms of cough or phlegm of different duration (1-2 months a year and 3 or more months a year). Light and severe asthmatic symptoms, as well as symptoms of cough or phlegm of different persistence, did not show a clear pattern with traffic indicators; only for truck traffic the data suggest a trend for stronger associations in subjects with more severe symptoms.

The relationship between these traffic indicators and the respiratory outcomes was explored in detail (Table 4) evaluating the associations with asthma symptoms without cough or phlegm, cough or phlegm without asthma and the combination of the two conditions. Asthma symptoms, when alone, were only weakly associated with high traffic density, continuous car and continuous truck transit. Traffic exposure indicators were more strongly associated with cough or phlegm, and particularly when accompanied by asthma symptoms. For this last condition (asthma with cough or phlegm) there were strong and significant associations for all the indicators.

The analyses of the independent and joint effects of truck traffic and car traffic on the risk of cough or phlegm are shown in Table 5. In some categories the numbers are relatively small and the effect

estimates are therefore unstable. The table confirms an independent effect of truck transit on cough or phlegm (OR=1.50; 95% CI: 0.93, 2.43), and also shows a similar association for car transit (OR=1.30; 95% CI: 0.94, 1.81).

Further analyses were performed to explore the possible effect modifiers of the association with frequent or continuous truck transit (Figure 1). Higher risks were observed for females for asthma symptoms (OR=1.34; 95% CI: 1.17, 1.53) and cough or phlegm (OR=1.62; 95% CI: 1.33, 1.98) than for males (OR=1.08; 95% CI: 0.94, 1.24; and OR=1.35; 95% CI: 1.14, 1.60 respectively). The pattern was present for both age groups. Cough or phlegm were more associated with truck transit in the Northern areas of the Country (OR=1.75; 95% CI: 1.47, 2.08) than in the Central and in the Southern areas (OR=1.37; 95% CI: 1.10, 1.71 and OR=1.09; 95% CI: 0.78, 1.51). Asthma symptoms were more associated with truck transit in the urban or rural areas (OR=1.31; 95% CI: 1.16, 1.47) than in the metropolitan areas (i.e. municipalities with more than 500.000 inhabitants) (OR=1.03; 95% CI: 0.87, 1.21). For both symptoms, higher associations (not statistically significant) were observed for the lowest level of parental education (13 years or less) than for the highest levels (high school/University). Asthma symptoms were more associated with truck transit when parental asthma or allergies were not reported. Age, smoking, presence of mould stain, change of residence and floor level did not act as effect modifiers. Comparable results were observed for the indicator of continuous car traffic (Figure 1 appendix).

Discussion

This study confirms our previously-reported associations between frequency and density of traffic in the street of residence and current respiratory symptoms in children [18]. For all indicators of traffic density and frequency, an increase in symptoms by level of exposure was observed. The strongest associations were for cough or phlegm and truck traffic, although an independent effect of

continuous car traffic was found. Traffic fume exposure was weakly associated with asthma symptoms, but the association was stronger if asthma was accompanied by cough or phlegm.

Few published studies have analyzed the effects of traffic in relationship to combinations of respiratory outcome such as asthma symptoms and persistent cough or phlegm. A positive association of air pollution with bronchitis symptoms, but not with asthma, has been reported in some studies [3, 6]. Other studies found both associations; e.g. Kim et al. (2004) found a significant increase in the risk of bronchitic symptoms and asthma in relation to higher level of traffic related to pollution in a US area with relatively clean air [22]. However, the findings of different studies should be compared with caution, because the definition of asthma adopted may be different and it can often include symptoms that may be related to bronchitis rather than to asthma [23]. In fact we have found an increased risk of asthma symptoms in children exposed to car and truck traffic fumes, essentially restricted to those who also reported cough or phlegm. There are two possible explanations for these findings. Firstly, it is possible that these children only had chronic bronchitis and did not have asthma symptoms, i.e. that they were “misdiagnosed” by our asthma symptom questionnaire [24]. The second, and perhaps more likely explanation, is that these children experienced an increased risk for a form of asthma that also involved persistent cough and phlegm.

Several previous questionnaire-based studies and surveys that used objective traffic counts as exposure metrics, have suggested that heavy traffic powered by diesel engines is more harmful for respiratory health than light traffic powered by gasoline engines [13, 23]. When we analyzed the effects of car and truck traffic separately on cough or phlegm, we found a stronger association with heavy traffic exposure (Table 4), consistent with the findings from other studies [4, 7, 9]. In the study conducted by Janssen and coworkers (2003), chronic respiratory symptoms were positively associated with high truck traffic and with air pollutants related to truck traffic counts [7]. In a cross-sectional study conducted on a sample of 1242 children (7-12 years old) [9], respiratory

symptoms were more prevalent among children attending schools located close (<100 m) to highways, particularly if the highway had a high truck traffic density.

We found significant associations of car traffic with cough or phlegm, although these were generally weaker than the associations for truck traffic (Tables 4 and 5). These findings could indicate that car traffic may also represent a source of pollutants increasing the risk of chronic cough or phlegm. However, these findings could also be related to the very high, and still increasing, number of cars powered by diesel engines at present circulating in Italy [25], and does not necessarily indicate that other types of car fume exposure also increase the risk of respiratory symptoms.

Children exposed to both, continuous car traffic and frequent or continuous truck transit, showed the highest risk of cough or phlegm (Table 5). The joint effect of car and truck exposure (OR=1.60) is approximately what we would expect under an additive model ($1+0.30+0.50 = 1.80$), given the independent effects of cars (OR=1.30) and trucks (OR=1.50). This is consistent with the hypothesis that truck and car traffic involve similar exposures (e.g. diesel fumes) so that their effects would be additive.

With respect to exposure to heavy traffic and related respiratory outcomes, we found stronger associations in females, for both age groups. An increased susceptibility of girls to air pollution has been reported by several investigators [4, 9, 10, 22] and was partially observed also in SIDRIA-1 [18]. A higher pollutant-to-lung volume ratio that results from larger airways relative to lung size in girls than in boys and enhanced cholinergic irritability could be considered as possible explanations [26, 27]. A gradient in the geographical distribution of the ORs for truck exposure and cough or phlegm is present in our data. This trend is consistent with the hypothesis that geo-climatic factors characterizing the Northern regions do favor accumulation of atmospheric pollutants. We do not

have a direct interpretation for the strong association (although not statically significant) between truck frequency and cough or phlegm in children of less educated parents, but it should be noted that the same type of effect modification by socioeconomic status has been seen in other studies of air pollution, regarding short term [28, 29] and long term effects [30]. Parental asthma or allergy did not act as modifiers of the association between cough or phlegm and truck traffic. For asthma symptoms, the association was stronger among parents that did not report asthma or allergies. These findings are reassuring with regards to the possibility of a reporting bias due to an over-reporting of traffic information among parents suffering from asthma or allergic disorders (in contrast with the suggestions of Piro *et al* 2008 [31]).

In contrast with the findings of the previous phase of SIDRIA, for symptoms of cough or phlegm we did not observe stronger associations among subjects living in metropolitan areas compared to those living in less urbanized areas; on the contrary, stronger associations were observed for asthma symptoms among urban-rural areas. A possible interpretation may be the presence of higher contrasts in pollution levels in urban-rural areas than in metropolitan areas.

The main strength of the present study is that it was based on a large random general population sample with a high response rate (86.7%). The information on symptoms of asthma and cough or phlegm included items derived from standardized validated questionnaires, that have been used in previous epidemiological studies in this population [32] and internationally [17]. The main weakness of our study is that it relies for both symptoms and exposure on self-reported information. Recently, the quality of self reported exposure has been questioned, particularly in studies on respiratory health in children [31, 33, 34]. The accuracy of the information about traffic exposure reported by parents has been examined in the first phase of SIDRIA. In a study in the city of Turin, the address of each respondent was linked to objective data on traffic available from a survey conducted by the Local Administration in 80 selected streets. The categories of the “truck frequencies” in the street of residence were found to be good predictors of the hourly median

number of heavy vehicles passing in the street; separate assessment for parents of symptomatic and non-symptomatic children showed no significant differences [18]. In the second phase of SIDRIA, we repeated the same analysis using objective data of traffic routinely collected, and, again, we found that the daily median number of vehicles was well predicted by the categories of traffic frequency, without significant differences between symptomatic and not symptomatic subjects (Migliore et al., manuscript in preparation). In a subset of the second phase SIDRIA sample we also performed several analyses in order to evaluate the existence of a possible response bias, using a method comparable to that used by Kuehni et al [33]. The results are presented in detail in the manuscript of Migliore et al. (in preparation). For the indicator of traffic density in the area of residence they do not support the existence of a response bias. For the indicators of truck or car traffic in the street of residence the results are less clearly interpretable, but the sub samples available for these matched analyses were very small and heavily selected, raising concerns regarding the generalizability and the interpretability of these analytical procedures.

The possibility of important residual confounding in our results is low, as we adjusted for several known or suspected risk factors for respiratory symptoms; however, we lacked some outdoor and indoor information, like ventilation characteristics of the house. Indeed, in Italy different times of windows' opening have been reported between North and Central Italy [35], even if the magnitude of the effect was small on indoor air pollutants concentrations.

Our questionnaire was focused on traffic pollution in the street and zone of residence, but did not collect information about other exposures such as those close to the schools. The absence of these details could have led to a small random misclassification of the exposures in the participants, while it seems unlikely as an explanation of the observed positive associations.

For reasons of consistency and comparability between age groups, we based the analyses on the parental questionnaires for both age-groups. However, in a sensitivity analysis we examined the associations between traffic indicators and respiratory symptoms including the answers on

symptoms given by adolescents (Table 4 appendix). The results for the outcome of cough or phlegm were substantially unchanged.

Conclusions

Although the specific set of airborne toxicants that facilitate and promote respiratory effects are still not known, our findings add further evidence to support a causal effect of exposure to traffic road pollution on respiratory illnesses in childhood. The adverse effects of air pollution represent a relevant public health problem, since exposure to traffic exhausts involves a large proportion of children having found that about 60% of children in our study are exposed to frequent or continuous car traffic and that about 20% of children are exposed to frequent or continuous truck traffic fumes.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

GB was involved in statistical analysis, interpretation of the results, and drafted the manuscript. EM assisted in the planning of the study, data collection, data management and analysis and helped drafting the manuscript. CG was national coordinator of the study, conceived the study and participated in its design and conduction and in the manuscript preparation. NP contributed to data analysis, interpretation of the results, and critical review of the manuscript. FF, LB, ECh, GG, PS, VDO, LA, CG and GC were local coordinator and were involved in the conception and design of the study, acquisition of local data, interpretation of the results, and critical review of the manuscript. AB and GV participated in the study design, statistical analyses, and revision of the manuscript. MB, RP and LA, as respiratory specialists, were involved in medical data collection, interpretation of the results and revision of the paper. EC, as air quality specialist, contributed to exposure assessment and critical review of the manuscript. All of the authors have read and approved the final version of the manuscript.

Acknowledgements:

The authors would like to thank the pupils and their parents and the staff of the schools for their co-operation. They specially thank Prof. Franco Merletti and Prof. Stefania La Grutta for valuable comments on the manuscript. This work was also supported by the Progetto Lagrange, Fondazione CRT / ISI, and the Health Research Council of New Zealand. The SIDRIA project was partially funded by the Italian Ministry of Health - Regional Health Agency, Bologna. This study was partly supported by the Piedmont Region. Enrica Migliore's work on this paper was partially supported by the Master in Epidemiology, University of Turin.

References

1. Behrens T, Taeger D, Maziak W, Duhme H, Rzehak P, Weiland SK, Keil U: **Self-reported traffic density and atopic disease in children. Results of the ISAAC Phase III survey in Muenster, Germany.** *Pediatr Allergy Immunol* 2004, **15**:331-339.
2. Brauer M, Hoek G, Van Vliet P, Meliefste K, Fischer PH, Wijga A, Koopman LP, Neijens HJ, Gerritsen J, Kerkhof M, Heinrich J, Bellander T, Brunekreef B: **Air pollution from traffic and the development of respiratory infections and asthmatic and allergic symptoms in children.** *Am J Respir Crit Care Med* 2002, **166**:1092-1098.
3. Braun-Fahrlander C, Vuille JC, Sennhauser FH, Neu U, Kunzle T, Grize L, Gassner M, Minder C, Schindler C, Varonier HS, Wuthrich B: **Respiratory health and long-term exposure to air pollutants in Swiss schoolchildren. SCARPOL Team. Swiss Study on Childhood Allergy and Respiratory Symptoms with Respect to Air Pollution, Climate and Pollen.** *Am J Respir Crit Care Med* 1997, **155**:1042-1049.
4. Brunekreef B, Janssen NA, de Hartog J, Harssema H, Knape M, van Vliet P: **Air pollution from truck traffic and lung function in children living near motorways.** *Epidemiology* 1997, **8**:298-303.
5. Duhme H, Weiland SK, Rudolph P, Wienke A, Kramer A, Keil U: **Asthma and allergies among children in West and East Germany: a comparison between Munster and Greifswald using the ISAAC phase I protocol. International Study of Asthma and Allergies in Childhood.** *Eur Respir J* 1998, **11**:840-847.
6. Hirsch T, Weiland SK, von Mutius E, Safeca AF, Grafe H, Csaplovics E, Duhme H, Keil U, Leupold W: **Inner city air pollution and respiratory health and atopy in children.** *Eur Respir J* 1999, **14**:669-677.
7. Janssen NA, Brunekreef B, van Vliet P, Aarts F, Meliefste K, Harssema H, Fischer P: **The relationship between air pollution from heavy traffic and allergic sensitization, bronchial hyperresponsiveness, and respiratory symptoms in Dutch schoolchildren.** *Environ Health Perspect* 2003, **111**:1512-1518.
8. Nicolai T, Carr D, Weiland SK, Duhme H, von Ehrenstein O, Wagner C, von Mutius E: **Urban traffic and pollutant exposure related to respiratory outcomes and atopy in a large sample of children.** *Eur Respir J* 2003, **21**:956-963.
9. van Vliet P, Knape M, de Hartog J, Janssen N, Harssema H, Brunekreef B: **Motor vehicle exhaust and chronic respiratory symptoms in children living near freeways.** *Environ Res* 1997, **74**:122-132.
10. Venn AJ, Lewis SA, Cooper M, Hubbard R, Britton J: **Living near a main road and the risk of wheezing illness in children.** *Am J Respir Crit Care Med* 2001, **164**:2177-2180.
11. Pandya RJ, Solomon G, Kinner A, Balmes JR: **Diesel exhaust and asthma: hypotheses and molecular mechanisms of action.** *Environ Health Perspect* 2002, **110 Suppl 1**:103-112.
12. Sydbom A, Blomberg A, Parnia S, Stenfors N, Sandstrom T, Dahlen SE: **Health effects of diesel exhaust emissions.** *Eur Respir J* 2001, **17**:733-746.
13. Binková B, Bobak M, Chatterjee A, Chauhan A, Dejmek J, Dockery D, Everard M, Forastiere F, Gilliland F, Holgate S, Johnston S, Krzyzanowski M, Kuna-Dibbert B, Maynard R, Raaschou-Nielsen O, Samet J, Schneider J, Skerrett P, Šrám R, Walters D, Weiland S, Winneke G: **WHO Monograph: The effects of air pollution on children's health and development: a review of the evidence.** *WHO Regional Office for Europe* 2004.
14. SIDRIA: **Asthma and respiratory symptoms in 6-7 yr old Italian children: gender, latitude, urbanization and socioeconomic factors. SIDRIA (Italian Studies on Respiratory Disorders in Childhood and the Environment).** *Eur Respir J* 1997, **10**:1780-1786.

15. Simoni M, Lombardi E, Berti G, Rusconi F, La Grutta S, Piffer S, Petronio MG, Galassi C, Forastiere F, Viegi G: **Mould/dampness exposure at home is associated with respiratory disorders in Italian children and adolescents: the SIDRIA-2 Study.** *Occup Environ Med* 2005, **62**:616-622.
16. Galassi C, De Sario M, Biggeri A, Bisanti L, Chellini E, Ciccone G, Petronio MG, Piffer S, Sestini P, Rusconi F, Viegi G, Forastiere F: **Changes in prevalence of asthma and allergies among children and adolescents in Italy: 1994-2002.** *Pediatrics* 2006, **117**:34-42.
17. Asher MI, Keil U, Anderson HR, Beasley R, Crane J, Martinez F, Mitchell EA, Pearce N, Sibbald B, Stewart AW, Strachan D, Weiland SK, Williams HC: **International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods.** *Eur Respir J* 1995, **8**:483-491.
18. Ciccone G, Forastiere F, Agabiti N, Biggeri A, Bisanti L, Chellini E, Corbo G, Dell'Orco V, Dalmaso P, Volante TF, Galassi C, Piffer S, Renzoni E, Rusconi F, Sestini P, Viegi G: **Road traffic and adverse respiratory effects in children. SIDRIA Collaborative Group.** *Occup Environ Med* 1998, **55**:771-778.
19. Ellwood P, Asher MI, Beasley R, Clayton TO, Stewart AW: **The international study of asthma and allergies in childhood (ISAAC): phase three rationale and methods.** *Int J Tuberc Lung Dis* 2005, **9**:10-16.
20. de Jongste JC, Shields MD: **Cough . 2: Chronic cough in children.** *Thorax* 2003, **58**:998-1003.
21. Galassi C, Forastiere F, Biggeri A, Gabellini C, De Sario M, Ciccone G, Biocca M, Bisanti L: **[SIDRIA second phase: objectives, study design and methods].** *Epidemiol Prev* 2005, **29**:9-13.
22. Kim JJ, Smorodinsky S, Lipsett M, Singer BC, Hodgson AT, Ostro B: **Traffic-related air pollution near busy roads: the East Bay Children's Respiratory Health Study.** *Am J Respir Crit Care Med* 2004, **170**:520-526.
23. Brunekreef B, Sunyer J: **Asthma, rhinitis and air pollution: is traffic to blame?** *Eur Respir J* 2003, **21**:913-915.
24. Seear M, Wensley D: **Chronic cough and wheeze in children: do they all have asthma?** *Eur Respir J* 1997, **10**:342-345.
25. ENEA: *Rapporto Energia e Ambiente 2005*. Roma: Ente per le Nuove Tecnologie, l'Energia e l'Ambiente; 2005.
26. Annesi-Maesano I, Agabiti N, Pistelli R, Couilliot MF, Forastiere F: **Subpopulations at increased risk of adverse health outcomes from air pollution.** *Eur Respir J Suppl* 2003, **40**:57s-63s.
27. Forastiere F, Corbo GM, Pistelli R, Michelozzi P, Agabiti N, Brancato G, Ciappi G, Perucci CA: **Bronchial responsiveness in children living in areas with different air pollution levels.** *Arch Environ Health* 1994, **49**:111-118.
28. Jerrett M, Burnett RT, Brook J, Kanaroglou P, Giovis C, Finkelstein N, Hutchison B: **Do socioeconomic characteristics modify the short term association between air pollution and mortality? Evidence from a zonal time series in Hamilton, Canada.** *J Epidemiol Community Health* 2004, **58**:31-40.
29. Martins MC, Fatigati FL, Vespoli TC, Martins LC, Pereira LA, Martins MA, Saldiva PH, Braga AL: **Influence of socioeconomic conditions on air pollution adverse health effects in elderly people: an analysis of six regions in Sao Paulo, Brazil.** *J Epidemiol Community Health* 2004, **58**:41-46.
30. Pope CA, 3rd, Burnett RT, Thun MJ, Calle EE, Krewski D, Ito K, Thurston GD: **Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution.** *JAMA* 2002, **287**:1132-1141.

31. Piro FN, Madsen C, Naess O, Nafstad P, Claussen B: **A comparison of self reported air pollution problems and GIS-modeled levels of air pollution in people with and without chronic diseases.** *Environ Health* 2008, **7**:9.
32. Fuso L, de Rosa M, Corbo GM, Valente S, Forastiere F, Agabiti N, Pistelli R: **Repeatability of the ISAAC video questionnaire and its accuracy against a clinical diagnosis of asthma.** *Respir Med* 2000, **94**:397-403.
33. Kuehni CE, Strippoli MP, Zwahlen M, Silverman M: **Association between reported exposure to road traffic and respiratory symptoms in children: evidence of bias.** *Int J Epidemiol* 2006, **35**:779-786.
34. Heinrich J, Gehring U, Cyrus J, Brauer M, Hoek G, Fischer P, Bellander T, Brunekreef B: **Exposure to traffic related air pollutants: self reported traffic intensity versus GIS modelled exposure.** *Occup Environ Med* 2005, **62**:517-523.
35. Simoni M, Biavati P, Carrozzi L, Viegi G, Paoletti P, Matteucci G, Ziliani G, Ioannilli E, Sapigni T: **The Po river Delta (North Italy) Indoor Epidemiological Study: home characteristics, indoor pollutants, and subjects' daily activity pattern.** *Indoor Air* 1998, **8**:70-79.

Figure legends

Figure 1: Associations between exposure to truck transit and respiratory symptoms, by several characteristics. The figure reports the associations (OR and 95% CI) between exposure to truck transit (frequent/continuous vs never) and asthma symptoms and cough or phlegm, by different factors (age, gender, latitude, level of urbanization, parental education, indoor mould/dampness, change of residence, floor of the apartment, passive smoke at home, parental asthma or allergies). All ORs were adjusted for potential confounder, excluding the stratification factor. Statistical significance (p values) of the interaction terms are reported.

Tables

Table 1. Prevalence of respiratory outcomes investigated in a sample of Italian schoolchildren.

		Asthma symptoms		
		YES	NO	TOTAL
		n (%)	n (%)	
Cough or phlegm	YES			
	n (%)	961 (2.9%)	1337 (4.0%)	2298 (6.8%)
	NO			
	n (%)	3591 (10.7%)	27743 (82.5%)	31334 (93.2%)
TOTAL		4552 (13.5%)	29080 (86.5%)	33632 (100%)

Table 2. Prevalence (%) of current respiratory symptoms by various subgroups (N=33632)

	Asthma symptoms	Cough or phlegm
	% (n)	% (n)
Total	13.5 (4552)	6.8 (2298)
Age		
6-7 year olds	13.7 (2736)	8.4 (1687)
13-14 year olds	13.3 (1816)	4.5 (611)
Gender		
male	14.6 (2531)	7.4 (1275)
female	12.4 (2012)	6.3 (1019)
Parental asthma or allergies		
yes	17.6 (2376)	8.9 (1202)
no	10.8 (2176)	5.4 (1096)
Urbanization level		
<10,000 inhabitants	11.2 (623)	5.3 (295)
10-100,000 inhabitants	13.2 (1246)	5.6 (534)
100-500,000 inhabitants	14.8 (973)	7.1 (468)
>500,000 inhabitants	14.2 (1710)	8.3 (1001)
Exposure to passive smoke		
yes	14.8 (2509)	7.4 (1254)
no	12.2 (2019)	6.2 (1030)
Presence of mould/dampness		
yes	17.4 (631)	8.7 (314)
no	12.9 (3598)	6.5 (1807)
<i>missing</i>	<i>14.8 (323)</i>	<i>8.1 (177)</i>
Parental education		
primary school	15.1 (207)	7.8 (107)
secondary school	14.2 (1420)	7.4 (734)
high school	13.3 (2031)	6.5 (990)
university	12.7 (853)	6.6 (443)
Floor level of apartment		
Ground	13.5 (1273)	6.2 (585)
1 st -2 nd floor	13.9 (1962)	6.6 (933)
3 rd floor or more	12.9 (1231)	7.7 (738)
Change of residence		
Yes	12.6 (2182)	6.3 (1099)
No	14.5 (2174)	7.4 (1109)

Table 3. Associations between traffic indicators and asthma symptoms and cough or phlegm

	Asthma symptoms				Cough or phlegm			
	n cases	(%)	OR*	95% CI	n cases	(%)	OR*	95% CI
Traffic density								
Absent	564	12.5	1.00		270	6.0	1.00	
Low	1228	12.6	1.04	0.93-1.16	558	5.7	0.94	0.80-1.10
Moderate	1375	13.8	1.13	1.01-1.26	689	6.9	1.04	0.89-1.22
High	795	14.6	1.17	1.03-1.33	475	8.7	1.24	1.04-1.49
Frequency cars transit								
Never	309	12.7	1.00		142	5.9	1.00	
Sometimes	1147	12.1	0.95	0.83-1.10	517	5.5	0.90	0.74-1.10
Frequently	1399	13.6	1.06	0.93-1.22	679	6.6	1.02	0.84-1.26
Continuously	1108	14.9	1.16	1.00-1.33	656	8.8	1.32	1.08-1.63
Frequency trucks transit								
Never	1412	12.3	1.00		652	5.7	1.00	
Sometimes	1644	13.6	1.10	1.02-1.19	812	6.7	1.14	1.02-1.26
Frequently	682	14.5	1.16	1.05-1.29	399	8.5	1.41	1.23-1.61
Continuously	211	15.7	1.27	1.08-1.50	132	9.8	1.67	1.36-2.06

* All ORs were adjusted for study centre, age, sex, parental asthma or allergies, parental education, passive smoke at home, indoor moulds, floor level of the apartment and change of residence.

Table 4. Associations between traffic indicators and asthma symptoms with or without cough or phlegm.


	Asthma symptoms WITHOUT cough or phlegm				Cough or phlegm WITHOUT asthma symptoms				Asthma symptoms WITH cough or phlegm			
	n	(%)	OR*	95% CI	n	(%)	OR*	95% CI	n	(%)	OR*	95% CI
Traffic density												
Absent	465	10.3	1.00		171	3.8	1.00		99	2.2	1.00	
Low	996	10.2	1.00	0.89-1.13	326	3.4	0.83	0.69-1.00	232	2.4	1.12	0.88-1.42
Moderate	1083	10.9	1.08	0.95-1.22	397	4.0	0.92	0.75-1.11	292	2.9	1.27	1.02-1.57
High	601	11.0	1.13	0.99-1.28	281	5.2	1.14	0.92-1.41	194	3.6	1.52	1.17-1.96
Frequency cars transit												
Never	254	10.5	1.00		87	3.6	1.00		55	2.3	1.00	
Sometimes	938	9.9	0.93	0.80-1.07	308	3.3	0.91	0.72-1.15	209	2.2	0.97	0.71-1.34
Frequently	1129	11.0	1.03	0.89-1.19	409	4.0	1.07	0.83-1.36	270	2.6	1.10	0.80-1.52
Continuously	823	11.0	1.07	0.92-1.25	371	5.0	1.30	1.02-1.67	285	3.8	1.53	1.11-2.10
Frequency trucks transit												
Never	1158	10.1	1.00		398	3.5	1.00		254	2.2	1.00	
Sometimes	1299	10.8	1.07	0.98-1.17	467	3.9	1.10	0.95-1.26	345	2.9	1.27	1.09-1.48
Frequently	525	11.1	1.13	1.01-1.26	242	5.1	1.44	1.23-1.69	157	3.3	1.51	1.21-1.87
Continuously	150	11.2	1.21	1.01-1.46	71	5.3	1.47	1.13-1.93	61	4.5	2.11	1.58-2.81

* All ORs were adjusted for study centre, age, sex, parental asthma or allergies, parental education, passive smoke at home, indoor moulds, floor level of the apartment and change of residence.

Table 5. Associations between combined exposure to truck and car transit and cough or phlegm.

		Trucks transit			
		Never	Sometimes	Freq/continuously	
Cars transit	Never/ sometimes	<i>n cases (%)</i> OR* (95% CI)	426 (5.3) 1.00	211 (5.9) 1.12 (0.95-1.32)	19 (8.2) 1.50 (0.93-2.43)
	Frequently	<i>n cases (%)</i> OR* (95% CI)	168 (6.3) 1.09 (0.90-1.33)	370 (6.4) 1.12 (0.96-1.30)	140 (7.7) 1.42 (1.15-1.75)
	Continuously	<i>n cases (%)</i> OR* (95% CI)	56 (7.8) 1.30 (0.94-1.81)	228 (8.4) 1.42 (1.19-1.70)	372 (9.3) 1.60 (1.36-1.87)

* All ORs were adjusted for presence of current asthma symptoms, study centre, age, sex, parental asthma or allergies, parental education, passive smoke at home, indoor moulds, floor level of the apartment and change of residence.



Additional files provided with this submission:

Additional file 1: migliore_appendix ms.doc, 177K

<http://www.ehjournal.net/imedia/9721784882396933/supp1.doc>

Additional file 2: confidential draft paper migliore.doc, 573K

<http://www.ehjournal.net/imedia/2400280252396924/supp2.doc>