

Selenium may be adversely associated with kidney function and kidney length in adult offspring of parents with Balkan Endemic Nephropathy and controls.

A two-year follow-up study

Wilfried Karmaus ^{1#}, Plamen Dimitrov ², Valeri Simeonov ³, Rossitza Georgieva ², Angel Bonev ³, Svetla Tsolova ²

¹ Department of Epidemiology and Biostatistics, University of South Carolina, Columbia, South Carolina 29208, United State of America

² National Center of Public Health Protection, Sofia, Bulgaria

³ Vratza District Hospital, Vratza, Bulgaria

E-mail addresses: Wilfried Karmaus *- karmaus@sc.edu; Plamen Dimitrov - p.dimitrov@ncphp.government.bg; Valeri Simeonov - p.dimitrov@ncphp.government.bg; Rossitza Georgieva - r.georgieva@ncphph.government.bg; Angel Bonev - p.dimitrov@ncphp.government.bg; Svetla Tsolova - sv.tzolova@ncphp.government.bg

* Corresponding author

Abstract

Background

There is concern that exposures to metals (cadmium and lead) and metalloids (arsenic and selenium) are related Balkan Endemic Nephropathy (BEN), a tubulointerstitial kidney disease.

Methods

In 2003/04 we recruited 102 adults whose parents had BEN and who resided in one of three communities (Vratza, Bistretz, or Beli Izvor, Bulgaria). A control group comprised of 99 adults having non-BEN hospitalized parents was enrolled in the study during the same time. We conducted face-to-face interviews, ultrasound kidney measurements, and determined kidney function in two consecutive investigations (2003/04 and 2004/05). Metals and metalloids were measured in urine and blood samples. To assess the agreement between these consecutive measurements, we calculated intraclass correlation coefficients. Repeated measurement data were analyzed using mixed models.

Results

We found that cadmium and arsenic were associated with neither kidney size nor function. Lead had a significant but negligible effect on creatinine clearance. Selenium showed a weak but significant adverse association with three of the four kidney parameters, namely kidney length, creatinine clearance, and β_2 -microglobulin. These associations were not restricted to the offspring of BEN patients. Adding credence to these findings are reports showing comparable adverse kidney effects in animals exposed to selenium.

Conclusions

The findings of this 2-year follow-up study indicate the importance of exploring further whether selenium results in adverse renal effects in humans.

Background

Balkan Endemic Nephropathy (BEN) is a tubulointerstitial kidney disease that progresses slowly over many years. First characterized in the Vratza District, Bulgaria in 1956 [1], similar nephropathies were subsequently described in Yugoslavia [2, 3] and in Romania [4]. In 1964 this disease was recognized as a new nosological entity and named Balkan Endemic Nephropathy. The final disease stage is characterized by renal failure and shrinkage of both kidneys to the size of walnuts [5]. BEN shows familial clustering and develops only in certain areas in the Balkan countries – Bulgaria, Romania, Serbia, Montenegro, Croatia, and Bosnia and Herzegovina. Its spatial distribution shows a mosaic-like pattern, with some villages afflicted for decades, whereas others situated in the same vicinity have remained free of BEN [6-8]. It is still unknown what causes BEN. Some studies claim that exposures to metals (cadmium and lead) and metalloids (arsenic and selenium) are related to the disease, with selenium being protective [9-12].

An important characteristic of BEN is the late onset of the disease (not before age 50) and thus probably a long latency period and/or a period of being at risk between exposure and the gradually developing clinical manifestation of the disease. To determine exposures before the onset of the disease, we chose to study the offspring of

patients with BEN, who are at a higher risk for the disease, and use the offspring of other non-BEN patients as a control group.

Early markers for BEN are a lower creatinine clearance, increased excretion in urine of total protein, albumin, and β_2 -microglobulin, and symmetrically decreased kidney measures [13-24]. We recently reported in a cross-sectional investigation that changes in these markers are associated with being offspring of BEN parents [25]. In this work we link clinical markers with the concurrent burden of metals and metalloids.

We investigated four metals and metalloids: lead, arsenic, cadmium, and selenium.

Lead, absorbed in blood, is bound extensively to erythrocytes (90-95%) and lead content in whole blood is used routinely as a biomarker of exposure (up to a few months before sampling) [26, 27]. Most arsenic absorbed by the lungs or by the gastro-intestinal tract is excreted in the urine, mainly within 1-2 days. Measurement of urinary arsenic levels is generally accepted as the most reliable indicator of recent exposure [28]. Total arsenic in urine, which is measured in this study, includes inorganic arsenic, as well as its metabolites monomethylarsonic acid and dimethylarsinic acid.

Cadmium is a heavy metal which accumulates predominantly in the kidneys (half-life: 15-40 years) and liver and is excreted mainly in urine. In low-level exposures (below 10 $\mu\text{g/g}$ creatinine) cadmium content in urine reflects body burden and is used as a biomarker of cumulative exposure [27, 29]. After absorption, selenium in blood is distributed between red blood cells and plasma/serum. The concentration in plasma or

serum is considered to reflect short-term exposure, whereas the selenium content of erythrocytes indicates long-term exposure [29]. In a steady state, the Se levels in red blood cells are considered to be higher than those in serum [30, 31], however, both are significantly correlated and comparably associated with daily intake [32-34]. Serum selenium is used for assessment of occupational and environmental selenium exposure, including nutritional status, in various studies, for instance, the U.S. National Health and Nutrition Examination Survey [29, 35, 36].

Based on prior reports in the BEN literature we formulated five hypotheses:

- 1) Selenium serum concentration is protective against adverse kidney effects.
- 2) Lead in blood, 3) cadmium in urine, and 4) arsenic in urine are associated with adverse alterations in BEN kidney markers and size.
- 5) Adult offspring with a parental history of BEN show stronger associations between metals and/or arsenic and kidney variables.

Methods

Study Population

From October, 2003, to April, 2004, we recruited 102 adults whose father and/or mother were registered in the Vratza Hospital as BEN patients and who resided in one of three communities (Vratza, Bistretz, or Beli Izvor, Bulgaria). The parental diagnosis of BEN was based on epidemiological, clinical/laboratory, and pathological/anatomical criteria [37]. Regarding epidemiology, the region is known to have a high prevalence of BEN. The individual criterion is onset after the age of 50. Clinical and laboratory criteria are as follows: obscure onset, chronic course, absence of edema, and normochromic anemia

in the more advanced phases. Pathologically, an almost symmetrical macroscopic shrinkage of the two kidneys is found. A control group of nearly equal size, 99 adult offspring of non-BEN hospitalized patients was enrolled at the same time. Diagnoses in control parents included diabetes mellitus, cardiovascular disorders, and liver problems. Only three of the 99 control subjects had parents with kidney diseases (one paternal kidney cancer not related to BEN and two maternal pyelonephritis cases). Subjects of both groups were frequency-matched according to gender and ten-year age groups. The population was enrolled and examined in 2003/04 and re-examined in 2004/05, which provided us with two repeated measurements one year apart.

All participants provided written consent through a procedure approved by the Institutional Review Board (human subject research committee) of the National Center of Public Health Protection, Sofia, Bulgaria. This research was reviewed and approved by the Commission of Medical Ethics at the National Center of Hygiene, Medical Ecology and Nutrition, Sofia, Bulgaria, on December 16, 2002.

Interviews

We conducted face-to-face interviews with all participants, either in the hospital or by visiting them in their home villages. The standardized questionnaire asked for the place of residence, type of water supply, diet, smoking and drinking habits, medical symptoms, family history of BEN, family history of other kidney diseases and kidney tumors, and occupational history.

Physical examination and determination of kidney sizes

Physical examinations, performed by an experienced physician with board certification in internal diseases and nephrology, assessed the general health status of the study subjects, including height and weight, and looked for symptoms of BEN and/or other internal diseases. Blood pressure was measured according to standards set by the World Health Organization [38].

Ultrasound examination of both kidneys included the patient lying on the left side and then on the right side, and took 20–30 minutes. The longest dimension of the kidney, the thickness of the kidney parenchyma (in the thinnest or minimal part), and the relationship of parenchyma and pyelon were measured. Data gathered also included parenchyma structure and the location, size, and morphology (cysts, stones, and tumors) of the kidneys. The images were saved electronically for future reference. The ultrasonographer (AGB) worked in the Department of Image Diagnostics, Vratza District Hospital, and was blinded to the clinical status of the participant (BEN or control offspring).

Urine and blood collection and determination of kidney markers

Prior to the examination day, study subjects were provided with 100 mL polypropylene containers and instructed to collect their urine, avoiding external contamination, immediately after rising in the morning and to take it to the laboratory without delay. They were also told to refrain from eating fish, seafood, chicken, and organ meat during the week before sample collection in order to diminish their intake of organic arsenic via

diet. At the laboratory, aliquots of the first morning urine sample were transferred to polypropylene tubes: one aliquot was used for the measurement of creatinine; the other two were acidified with either 1N HCl (Merck) for arsenic determination or 1N HNO₃ (Merck) for cadmium determination. Then the aliquots were frozen at -20°C.

After the morning urine collection, the participants underwent a 4-hour examination with urine and blood sampling. Study subjects were asked to empty their bladders and were given 0.5 L of water to drink. From that moment on, all their urine was collected in polyethylene containers and refrigerated: At the end of the 1st hour and at 4 hours the study subjects were asked to urinate. Venous blood was collected after 2 hours.

The hour-one urine was checked with pH-indicator paper strips (Merck). If the pH was lower than 7, a portion of the urine was alkalized by adding 1N NaOH until a pH equal to 7 or between 7 and 8 was reached. Creatinine and β_2 -microglobulin were measured in aliquots of the hour-one urine sample. The volume of the hour-one urine sample was taken into account when the total urine volume was recorded at the end of the 4-hour period. The total 4-hour urine sample was apportioned into aliquots and frozen at -20°C. As described elsewhere [25], we determined total protein, albumin, and creatinine in urine in the 4-hour sample. To adjust for creatinine concentration, it was measured three times: in the first morning, the hour-one, and the 4-hour samples.

Venous blood for hematology, the determination of lead, and for serum separation was taken in the middle of the 4-hour urine collection period. Blood samples for

determination of lead were drawn in K₂EDTA vacutainers. Serum was separated for the analyses of selenium and creatinine.

Creatinine clearance

Creatinine clearance (CCR) was estimated from serum creatinine values by using the Cockcroft and Gault (1976) formula. Specifically,

$$\text{CCR (ml/min)} = \frac{(140 - \text{age}) \times \text{body weight in kg}}{(\text{serum creatinine} \times 72)} \quad (\times 0.85 \text{ for females})$$

Analyses of trace elements

Lead, cadmium, arsenic, and selenium were determined using Perkin-Elmer atomic absorption spectrometers (Bodenseewerk Perkin-Elmer, Ueberlingen, Germany). A Model 4110 ZL atomic absorption spectrometer (AAS) with a transverse heated graphite atomizer (THGA) and longitudinal Zeeman-effect background correction equipped with an AS-72 autosampler, electrodeless discharge lamps (EDL) System II, software WinLab (Version 1.2) and both “Standard” (Part No B 300-0643) and “End-capped” (Part No B 300-0644) THGA tubes with integrated platforms was used for direct electro-thermal AAS measurements of lead in blood, selenium in serum, and cadmium in urine. A Model 3030 AAS with a Mercury/Hydride System MHS-20, automatic deuterium background corrector and EDL System II was used for direct hydride generation AAS determinations of arsenic (As) in urine.

To measure lead (Pb), blood was diluted 10-fold with 0.25% v/v Triton X-100; 10 µL of the sample and 25 µg of (NH₄)₂HPO₄ as a chemical modifier were injected into the

THGA, pre-treated with 250 µg of zirconium and 20 µg of iridium [39]. For selenium (Se), serum was diluted 5-fold with 0.1% v/v Triton X-100 in 0.2% HNO₃; 10 µL injections of the sample and 10 µg of rhodium modifier were injected into the THGA; standard addition calibration was applied [40]. For cadmium (Cd) in urine, the samples were diluted 5-fold with 0.05 v/v Triton X-100 in 0.2% HNO₃; 10 µL of the sample and 25 µg of (NH₄)₂HPO₄ as a chemical modifier were injected into the THGA, pre-treated with 250 µg of zirconium and 20 µg of iridium [39]. Matrix-matched calibrations were applied for Pb, Se, and Cd. Total As was determined after transformation of the inorganic arsenic, As(III) + As(V), monomethylarsonate and dimethylarsinate, into similarly reactive species by treatment with L-cysteine. The final dilution of the urine samples was 5-fold. 10 mL aliquots were directly analyzed by hydride generation AAS [41].

We used the following reference materials for trace elements: Lyphochek Urine Metals Control Level 1, Cat. No. 400 (Bio-Rad Laboratories, Germany); Seronorm Trace Elements Urine Blank Cat. No. 201305; Seronorm Trace Elements Serum Level 1, Cat. No. 201405, and Level 2, Cat. No. 203105; Seronorm Trace Elements Whole Blood, Level 1, Cat. No. 201505, and Level 2, Cat. No. 201605 (Sero AS, Norway).

Statistical analyses

Outcome variables were total protein, albumin, β₂-microglobulin, creatinine in serum, CCR, kidney cortex width, and kidney length. To answer the question whether these outcomes are related to selenium serum concentration, lead in blood, and cadmium and arsenic urine excretion, we controlled for potentially confounding factors such as

gender, age, history of smoking, diabetes mellitus, hypertension, body surface area (BSA), and creatinine in the morning urine sample, the 1-hour, and the 4-hour urine samples. For smoking, we distinguished current smoker, ex-smoker, and non-smoker. We asked for presence of diabetes mellitus and determined moderate hypertension in the first investigation (2003/04) and for new occurrence in the second investigation. BSA was used in the model to control for the effect of weight and height on kidney markers (Table 4). In each investigation (2003/04 and 2004/05) the average kidney cortex width and kidney length of both kidneys were used as predictors.

In order to estimate the association between repeated measurements of arsenic, cadmium, lead, selenium, and kidney markers, we applied linear mixed models. Measurements of the two investigations are not independent and mixed models allowed for adjusting for within-participant effects [42]. The mixed model assumes that the random effects and the error vector are normally distributed, which was the case for kidney sizes and CCR; β 2-microglobulin was log-transformed. SAS PROC MIXED was used to perform the regression analysis [43]. We used the Akaike information criteria and likelihood ratio tests to examine the significance of serial correlation (repeated statement) as well as to model random effects, along with a suitable variance-covariance matrix structure. To investigate whether offspring with a parental history of Balkan Endemic Nephropathy are more susceptible to metals and metalloids, we additionally stratified the models by parental history.

To compare the agreement between measurements of continuous variables in 2003/04

and 2004/05 we used the intra-class correlation coefficient (ICC). This is the between-subject minus the within-subject variance divided by the sum of the two variances. The ICC quantifies the proportion of total outcome variance that is due to inter-individual variation. A positive value, for instance, of 0.60 indicates that 60% of the variation is due to between-subject and 40% due to within-subject variance. ICC can become negative when the within-subject variance exceeds the between-subject variance. Due to methodological considerations [44], we chose not to directly adjust for creatinine using creatinine-adjusted parameters but to introduce creatinine as a confounder in the regression models.

Results

Of the 201 participants from the first year, 189 participated in the follow-up (94%). Two were deceased, one moved out of the area, two could not be contacted, and 7 decided not to participate further. Of the twelve lost, nine were offspring of BEN patients. With the exception of hypertension, potential risk factors show similar distributions in each of the two investigations (Table 1). Of the BEN offspring, 30.4% had moderate hypertension in 2003/04 and 29% in 2004/05. A lower prevalence was detected in control offspring (5.1 and 10.5%, respectively). BEN patients also reported more often a history of kidney stones and pyelonephritis.

Regarding arsenic and cadmium (measured in morning urine) and creatinine (measured in morning urine samples), we found no agreement between the two investigations for creatinine, a moderate agreement for arsenic and a good agreement for cadmium

(Spearman rank correlation and ICC). For creatinine in morning urine samples the within-subject variance exceeds the between-subject variance. When we used the directly creatinine-adjusted parameters (division by the creatinine concentrations) the agreement for cadmium and arsenic was reduced, simply because of the high intra-individual variance of creatinine. There was a better agreement for two measurements, namely lead in blood and selenium in serum, measured one year apart.

Regarding the kidney parameters, Spearman rank correlation and ICC were low for total protein and albumin in serum, for creatinine in the 1-hour and 4-hour urine sample, and for the glomerular filtration rate (Table 3). These values did not improve when we directly adjusted for creatinine, since the within-subject variance of creatinine was high. The agreement between the sonographic determinations of kidney dimensions (kidney cortex and kidney length), β_2 -microglobulin, and the estimated creatinine clearance (CCR) was sufficient to conclude that they measure some underlying disease processes. In the following analyses we focus on these four parameters to test whether arsenic and metals explain their variance.

We did not find any significant effect of arsenic and metals on kidney cortex width (Table 4). Of the metals and metalloids analyzed, only selenium showed an association with kidney length, β_2 -microglobulin, and the estimated CCR in the models presented in Table 4. Lead gained importance only in the control offspring. Similar to selenium, it seems to decrease CCR.

Higher selenium levels are associated with a longer kidney length in the total group and the BEN offspring. When comparing an individual with 70 µg/L with one having 20 µg/L selenium, on the average kidney length would be increased by 1 mm ($50 * 0.02$ mm). This is only a small increase compared to the total kidney length (approximately 120 mm). However, in BEN offspring the respective increase is 2 mm ($50 * 0.04$ mm; Table 4). Selenium seems to decrease CCR in the total group and the control group, but not in BEN offspring. When comparing an individual with 70 µg/L with one having 20 µg/L selenium, CCR would be reduced by 1.05 mL/min ($50 * 0.21$ mL) in the total group. This is a minimal decrease of CCR (range: 62-187 mL/min). In the control group, the decrease associated with lead is even smaller (0.45 mL/min). β_2 -microglobulin was log-transformed to achieve a normal distribution. A 50 µg/L increase in selenium would be approximately equivalent to a 1.4% increase in β_2 -microglobuline [$10^{(50 * 0.003)}$].

Discussion

Of elements investigated, cadmium and arsenic were associated neither with kidney dimension nor with function (hypotheses 3 and 4). Lead had a negligible effect on CCR (hypothesis 5). Selenium showed weak but significant associations with three of the four kidney parameters tested, namely kidney length, estimated CCR, and β_2 -microglobulin, but not kidney cortex width.

Our study included only BEN villages. Others think that the soil geochemistry in villages affected and not affected by BEN may be an etiologic factor [45]. It has been suggested that arsenic, cadmium, and lead (Pb) are elevated in the soil of BEN areas, whereas

selenium is deficient [9, 45, 46]. Ecological studies do not take individual exposures into account. Nevertheless, according to these investigations, we would have expected higher levels of arsenic, cadmium, and lead in our participants. To examine whether the levels in our samples were different, we compared these to other reports on metal and metalloid exposures in Bulgaria. Tsoleva *et al.*, when studying copper-smelter workers, described urine arsenic concentration of 7.73 μ g/L in a control group.[47] We found approximately 3 μ g/L in an unexposed population (Table 2). Regarding cadmium, two Bulgarian investigations reported urine levels of 0.8-1.0 μ g/L [48, 49]; we had urine concentrations of 0.6-0.7 μ g/L. However, adjusted for creatinine, urinary cadmium levels were higher in our Bulgarian samples (approximately 0.85 μ g/g creatinine) than levels reported for the United States (0.26 μ g/g creatinine)[50]. Lead blood concentration in our samples was lower (median of 90 μ g/L in 2003/4, Table 3) than levels reported for occupational exposures in Bulgaria [49, 51]. Selenium levels in this study (57-72 μ g/L, Table 2) were comparable to levels reported in other studies in Bulgaria (57 and 66.5 μ g/L; [27, 52]). Thus, concentrations detected in our samples were either comparable to or lower than levels in samples taken from other subpopulations in Bulgaria. Hence, these comparisons do not support the suggestion that BEN families are exposed to levels of arsenic, cadmium, selenium, or lead that could contribute to the occurrence of the disease.

Our study subjects were offspring of BEN patients who are likely to have a higher susceptibility of their kidneys to adverse exposures (hypothesis 5). Therefore, we expected to find that some of the elements would be associated with kidney function in

BEN offspring. However, compared to controls, BEN offspring were not more affected (Table 4). For lead (Pb) we found a weak association with creatinine clearance, but only in control offspring. Further, since our investigation is based on offspring who have not yet developed the disease, we believe that this excluded behavioral changes such as avoidance of chemicals, which could have distorted the results. Therefore, since prior studies that reported associations between arsenic/metals and BEN had an ecological design or were based on geochemistry analyses, our negative findings suggest that non-epidemiologic studies may have been misleading.

A limitation of our investigation is the high intra-individual variance of kidney parameters. If our analysis would have been based on a single examination, this could have led to biased results. However, we included repeated measurements that took intra-individual variances into account and provided added confidence in our results.

We are not aware of another study that investigated the association between kidney parameters and selenium in humans. Nevertheless, a recent study on selenium supplementation found an increased risk for type 2 diabetes.[53] We did not expect that selenium would have an adverse effect and be associated with increased β_2 -microglobulin and kidney length and decreased CCR. This may be a chance finding. However, the US Agency for Toxic Substances and Disease Registry lists a number of comparable kidney effects in animals following exposure to selenium at levels several hundred times higher than normal human intake. [35] Effects include hydropic degeneration in sheep [54], increases in kidney weights in mice and ducks [55-57],

larger Bowman's capsules and minimal papilla degeneration of the kidneys in rats [55, 58], and glomerulonephritis and proximal convoluted tubule nephropathy in long-tailed macaques.[59]

Conclusions

Our results do not support the hypotheses that cadmium, lead, or arsenic pose a risk for the development of Balkan Endemic Nephropathy, nor that selenium is protective. We found a weak adverse effect of selenium on kidney function and size in BEN offspring but also in controls. This is the first study that has investigated whether selenium is associated with renal health. Should our findings be corroborated, then there is a need to regulate selenium intake from all sources, including widely promoted nutritional supplements.

Competing interests

The authors declare that they have no competing interests.

Abbreviations: arsenic (As), atomic absorption spectrometer (AAS), Balkan Endemic Nephropathy (BEN), body surface area (BSA), cadmium (Cd); creatinine clearance rate (CCR), electrodeless discharge lamps (EDL), intra-class correlation coefficient (ICC), lead (Pb), selenium (Se), transverse heated graphite atomizer (THGA)

Authors' contributions

PD and WK designed the study, analyzed the data, and prepared the manuscript. VS and AB conducted the clinical investigations and helped with the preparation of the paper and the clinical interpretation. RG led the measurements of the metals and metalloids and reviewed the manuscript. ST provided toxicological expertise, help with the implementation of the study, data preparation, analyses, and the writing of the manuscript. All authors have read and approved the final manuscript.

Acknowledgements

The research was supported by NIH Research Grant # R01 TW06192 funded by the Fogarty International Center and the National Institute of Environmental Health Sciences, National Institutes of Health, USA, awarded to the National Center of Public Health Protection, Sofia, Bulgaria. We thank Susan Davis for revising the manuscript.

References

1. Tanchev Y, Evstatiev, Z., Dorossiev, D., Pencheva, J., Zvetkov, G.: Studies on the nephritides in the District of Vratza. *Savremenna Medicina* 1956, 7:14-29.
2. Danilovic V, Djuriscic M, Mokranjac M, Stojimirovic B, Zivojinovic J, Stojakovic P: Porodicna oboljenja bubrega u selu Sopic izazvana hronicnom intoksikacijom olovom. *Srpski Arh Tcelok Lek* 1957, 85:1115-1125.
3. Danilovic V, Djuriscic M, Mokranjac M, Stojimirovic B, Zivojinovic J, Stojakovic P: Chronic nephritis caused by poisoning with lead via the digestive tract (flour). *Presse Med* 1957, 65:2039-2040.
4. Fortza N, Negoescu, M.: Nefrita cronica azotemia endo-epidemic. *Stud Cercet Med* 1961, 1:217-221.
5. Vukelic M, Sostaric, B., Belicza, M.: Pathomorphology of Balkan endemic nephropathy. *Food Chem Toxicol* 1992, 30:193-200.
6. Tanchev Y, Dorossiev, D.: The first clinical description of Balkan endemic nephropathy (1956) and its validity 35 years later. In *Mycotoxins, Endemic Nephropathy and Urinary Tract Tumours*. Edited by Castenagro M, Plestina R, Dirheimer G, Chernozemsky IN, Bartasch H. Lyon: International Agency for Research on Cancer; 1991: 21-28
7. Radovanovic Z: Epidemiological characteristics of Balkan endemic nephropathy in Eastern regions in Yugoslavia. In *Mycotoxins, Endemic Nephropathy and Urinary Tract Tumours*. Edited by Castenagro M, Plestina R, Dirheimer G, Chernozemsky IN, Bartasch H. Lyon: International Agency for Research on Cancer; 1991: 11-20

8. Ganev V, Petropoulos E: Balkan Endemic Nephropathy and its Putative Environmental and Genetic Causes: A Review. In *6th International Conference of Environmental Science and Technology; Pythagorion, Samos, Greece*. Edited by Lekkas T. University of Aegen; 1999: 11-21.
9. Maksimovic ZJ: Selenium deficiency and Balkan endemic nephropathy. *Kidney Int Suppl* 1991, 34:S12-14.
10. Mihailovic M, Lindberg, P., Jovanovic, I., Antic, D.: Selenium status of patients with Balkan endemic nephropathy. *Biol Trace Elem Res* 1992, 33:71-77.
11. Long DT, Icopini G, Ganev V, Petropoulos E, Havezov I, Voice T, Chou K, Spassov A, Stein A: Geochemistry of Bulgarian soils in villages affected and not affected by Balkan endemic nephropathy: a pilot study. *Int J Occup Med Environ Health* 2001, 14:193-196.
12. Nichifor E, Balea M, Rusu G, Melencu M, Ghiordanescu N, Cristescu I, Dovlete C, Sonoc S: Studies on the familial character of endemic Balkan nephropathy. Possible role of the toxic hydric factor in the determination of "familial agglomerations" in endemic Balkan nephropathy. *Med Interne* 1985, 23:229-237.
13. Arsenovic A, Bukvic, D., Trbojevic, S., Maric, I., Djukanovic, L.: Detection of renal dysfunctions in family members of patients with balkan endemic nephropathy. *American Journal of Nephrology* 2005, 25:50-54.
14. Mitic-Zlatkovic M, Cukuranovic, R., Lecic, N., Stefanovic, V.: Urinary creatinine excretion in children from families with Balkan endemic nephropathy:

- evidence for genetic predisposition to the disease. *Pathologie Biologie* 2000, 48:554-557.
15. Stefanovic V, Mitic-Zlatkovic, M., Cukuranovic, R., Vlahovic, P.: Increased Urinary Protein Excretion in Children from Families with Balkan Endemic Nephropathy. *Nephron Clinical Practice* 2003, 95:116-120.
 16. Stefanovic V, Cukuranovic, R, Mitic-Zlatkovic, M, Hall, PW: Increased urinary albumin excretion in children from families with Balkan nephropathy. *Pediatr Nephrol* 2002, 17:913-916.
 17. Djukanovic L, Bukvic, D., Maric, I.: Creatinine clearance and kidney size in Balkan endemic nephropathy patients. *Clinical Nephrology* 2004, 61:384-386.
 18. Hall P, Vasilevic, M.: Beta2-microglobulin excretion as an index of renal tubular disorders with special reference to endemic Balkan nephropathy. *J Lab Clin Med* 1973, 81:897-904.
 19. Karlsson F, Lenkei, R.: Urinary excretion of albumin and beta 2-microglobulin in a population from an area where Balkan nephropathy is endemic. *Scand J Clin Lab Invest* 1977, 37:169-173.
 20. Hrabar A, Aleraj, B., Ceovic, S., Cvoriscec, D., Vacca, C., Hall, PW 3rd: Beta 2-microglobulin studies in endemic Balkan nephropathy. *Kidney Int Suppl* 1991, 34:S38-40.
 21. Hrabar A, Aleraj, B., Ceovic, S., Cvoriscec, D., Vacca, C., Hall, P. W.: A fifteen year cohort based evaluation of beta 2-microglobulin as an early sign of Balkan endemic nephropathy. *Kidney Int Suppl* 1991, 34:S41-43.

22. Stefanovic V, Mitic-Zlatkovic, M., Cukuranovic, R., Miljkovic, P., Pavlovic, N. M., Vlahovic, P.: Beta 2-microglobulin in patients with Balkan nephropathy and in healthy members of their families. *Kidney Int Suppl* 1991, 34:S21-26.
23. Jevremovic I, Jankovic, S., Radovanovic, Z., et al.: Beta2-microglobulinemia in a population exposed to Balkan endemic nephropathy: Inferences from repeated cross-sectional studies. *Kidney Int Suppl* 1991, 34:S35-S37.
24. Schadurin G, Statius van Eps, L.: Beta2-microglobulin: its significance in the evaluation of renal function. *Kidney International* 1987, 32:635-641.
25. Dimitrov P, Tsoleva S, Georgieva R, Bozhilova D, Simeonov V, Bonev A, Karmaus W: Clinical markers in adult offspring of families with and without Balkan Endemic Nephropathy. *Kidney International* 2006, (in print).
26. Cannon DJ: Heavy metal analysis. In *Clinical laboratory medicine*. Edited by Tilton RC, Balows, A., Hohnadel, D.C., Reiss, R.F. St. Louis, MI: Mosby-Year Book, Inc.; 1992: 1207
27. Tsalev D: Atomic absorption spectrometry in occupational and environmental health practice, vol. 3. CRC Press, Inc., Boca Raton, Florida, pp. 349. 1995.
28. Toxicological profile for arsenic (Draft for public comment)
[\[http://www.atsdr.cdc.gov/toxprofiles/tp2.html\]](http://www.atsdr.cdc.gov/toxprofiles/tp2.html)
29. Lauwerys R, Hoet, P. : *Industrial chemical exposure: Guidelines for biological monitoring*. 2nd edn. Boca Raton, FL: Lewis Publishers; 1993.
30. Rossi LC, Clemente GF, Santaroni G: Mercury and selenium distribution in a defined area and in its population. *Arch Environ Health* 1976, 31:160-165.

31. Schroeder HA, Frost DV, Balassa JJ: Essential trace metals in man: selenium. *J Chronic Dis* 1970, 23:227-243.
32. Neve J, Hanocq M, Molle L: [Application to various biological samples of a new method for selenium determination (author's transl)]. *J Pharm Belg* 1980, 35:345-350.
33. Raghunath R, Tripathi RM, Mahapatra S, Sadasivan S: Selenium levels in biological matrices in adult population of Mumbai, India. *Sci Total Environ* 2002, 285:21-27.
34. Longnecker MP, Stram DO, Taylor PR, Levander OA, Howe M, Veillon C, McAdam PA, Patterson KY, Holden JM, Morris JS, et al: Use of selenium concentration in whole blood, serum, toenails, or urine as a surrogate measure of selenium intake. *Epidemiology* 1996, 7:384-390.
35. Agency for Toxic Substances and Disease Registry: Toxicological Profile for Selenium. (Agency for Toxic Substances and Disease Registry - Division of Toxicology/Toxicology Information Branch ed. Atlanta, GA; 2003.
36. National Health and Nutrition Examination Survey: Documentation, Codebook, and Frequencies. Laboratory Component: Erythrocyte Protoporphyrin and Selenium. Survey years: 2003-2004. 2007:1-12.
37. Danilovic V.: Dijagnoza endemske nefropatije. *Radovi - XLVIII Od med nauka Akadnauka i umj BiH, Sarajevo* 1973, 17:53-64.
38. WHO: World Health Organization-International Society of Hypertension guidelines for the management of hypertension.
<http://new.euromise.org/mgt/who1999/who1999.html>; 1999.

39. Tsalev D.L., L. Lampugnani, R. Georgieva, K. Chakarova, Jr. IP: Electrothermal atomic absorption spectrometric determination of cadmium and lead with stabilized phosphate deposited on permanently modified platforms. *Talanta* 2002, 58 (2):331-340.
40. Tsalev D.L., L. Lampugnani, A. D'Ulivo, I.I. Petrov Jr., R. Georgieva, K. Marcucci, Zamboni R: Electrothermal atomic absorption spectrometric determination of selenium in biological fluids with rhodium modifier compared with hydride generation atomic spectrometric techniques. *Microchemical Journal* 2001, 70 (2):103-113.
41. Georgieva R, Petrov, P., Lampugnani, L., Tsalev, D.: Atomic Absorption Spectrometric Determination of Arsenic in Urine. In *2nd Black Sea Basin Conference on Analytical Chemistry; Sile-Istanbul, Turkey, September 14-17, 2003*. Book of Abstracts, p.218; 2003
42. Little RC, Miliken G, Stroup W, Wolfinger R: *SAS System for mixed models*. Cary, NC.: SAS Institute Inc; 1996.
43. SAS-Institute: SAS/STAT Software. Cary, NC: SAS Institute Inc.; 2000.
44. Schisterman EF, Whitcomb BW, Louis GM, Louis TA: Lipid adjustment in the analysis of environmental contaminants and human health risks. *Environ Health Perspect* 2005, 113:853-857.
45. Long D, Icopini, G., Ganev, V., Petropoulos, E., Havezov, I., Voice, T., Chou, K., Spassov, A., Stein, A.: Geochemistry of Bulgarian soils in villages affected and not affected by Balkan endemic nephropathy: A pilot study.

- International Journal of Occupational Medicine and Environmental Health* 2001, 14:193-196.
46. Angelieva R, Mladenova S: Trace Elements in Soil, Foodplants and Drinking Water in the Endemic Nephropathy Region in Bulgaria. In *Proceedings of the 4th symposium on Endemic (Balkan) Nephropathy, pp 59-61; Nis, Yugoslavia*. Edited by Strahinjc S, Stefanovic V. Institute of Nephrology and Hemodialysis, Faculty of Medicine, University of Nis, Nis, Yugoslavia; 1979
 47. Tsoleva S, Zaprianov Z, Dimitrov TS, Georgieva R, Khinkova L, Petrov I, Nikolova L: [The assessment of arsenic exposure in copper smelting]. *Probl Khig* 1995, 20:128-138.
 48. Binh N: Possibilities for clarifying of the relationship between Lead and Cadmium contents in the environment and in biological materials from different groups of people. University of Sofia, 1990.
 49. Hadjeva J: Proof-pathological problems of the independent and combined action of lead and cadmium. Clinic for professional diseases, Medical Academy, University of Sofia, 1991.
 50. Department of Health and Human Services: Third National Report on Human Exposure to Environmental Chemicals. In *NCEH Pub No 05-0570*. Atlanta, Georgia: National Center for Environmental Health, Division of Laboratory Sciences; 2005.
 51. Vaglenov A, Creus A, Laltchev S, Petkova V, Pavlova S, Marcos R: Occupational exposure to lead and induction of genetic damage. *Environ Health Perspect* 2001, 109:295-298.

52. Tzatchev KD: Rationalization of the trace element study in the chemical laboratory of the Medical Academy. University of Sofia, Medical Academy; 1994.
53. Stranges S, Marshall JR, Natarajan R, Donahue RP, Trevisan M, Combs GF, Cappuccio FP, Ceriello A, Reid ME: Effects of Long-Term Selenium Supplementation on the Incidence of Type 2 Diabetes: A Randomized Trial. *Ann Intern Med* 2007.
54. Smyth JB, Wang JH, Barlow RM, Humphreys DJ, Robins M, Stodulski JB: Experimental acute selenium intoxication in lambs. *J Comp Pathol* 1990, 102:197-209.
55. National Toxicology Program: NTP technical report on toxicity studies of sodium selenate and sodium selenite administered in drinking water to F344/N rats and B6C3F1 mice. In *Toxicity Report Series Number 38 NIH Publication 94-3387* (National Toxicology Program ed. Bethesda, MD: National Institutes of Health; 1994).
56. Johnson VJ, Tsunoda M, Sharma RP: Increased production of proinflammatory cytokines by murine macrophages following oral exposure to sodium selenite but not to seleno-L-methionine. *Arch Environ Contam Toxicol* 2000, 39:243-250.
57. Albers PH, Green DE, Sanderson CJ: Diagnostic criteria for selenium toxicosis in aquatic birds: dietary exposure, tissue concentrations, and macroscopic effects. *J Wildl Dis* 1996, 32:468-485.

58. Eckhert CD, Lockwood MK, Hsu MH, Ho J, Kang R: Microvascular changes in rat glomeruli as a consequence of small differences in selenium exposure. *Exp Mol Pathol* 1992, 57:222-234.
59. Cukierski MJ, Willhite CC, Lasley BL, Hendrie TA, Book SA, Cox DN, Hendrickx AG: 30-day oral toxicity study of L-selenomethionine in female long-tailed macaques (*Macaca fascicularis*). *Fundam Appl Toxicol* 1989, 13:26-39.

Table 1. Characteristics of the Cohort of Adult Offspring of BEN Patients and Controls

		2003/04		2004/05	
		Offspring of BEN patients, n=102 (%)	Offspring of control patients, n=99 (%)	Offspring of BEN patients, n=93 (%)	Offspring of control patients, n=96 (%)
Gender	Men	50.0	47.5	48.8	47.9
Age	30-40	32.4	15.5	32.3	15.6
	41-55	43.1	57.6	43.0	58.3
	56 plus	24.5	27.3	24.7	26.0
Place of birth	BEN village	95.0	72.7	94.6	72.9
Smoking status	current smoker	39.2	34.4	38.7	35.4
	ex-smoker	22.6	12.1	21.5	12.5
	non-smoker	38.2	53.5	39.8	52.1
Diabetes		7.8	6.1	5.4	7.3
Moderate hyper- tension in the offspring ξ		30.4	19.2	32.3	22.9
Other urinary tract diseases	kidney cancer	2.0	0	2.2	0
	kidney stones	10.8	4.0	10.8	6.5
	Pyelonephritis	10.8	0	9.7	1
	Hydronephrosis	1	0	1	0
	Cystitis	3.9	5.1	5.4	5.2
BSA $\#$	mean (standard deviation)	1.87 (0.22)	1.82 (0.21)	1.86 (0.18)	1.81 (0.17)

ξ Systolic blood pressure ≥ 140 and/or diastolic ≥ 90 mm Hg.

$\#$ BSA = square root ((height in cm \times weight in kg)/3600).

The intraclass correlation coefficient (ICC) is the between- subject minus the within-subject variance as a ratio to the sum of the two variances. A positive value of, for instance, 0.60 indicates that 60% of the variation is due to between-subject and 40% due to within-subject variance. ICC can become negative when the within-subject variance exceeds the between-subject variance.

Table 2. Metals and metalloids in urine and serum in two consecutive investigations

Variable		N	Median	5%- value	95%- value	Spearman correlation coefficient (p)	ICC	ICC 5%-value
As in urine (µg/L)	2003/04	201	3.10	0.70	9.30	0.28	-0.06	-0.18
	2004/05	189	2.90	0.80	8.90	(0.0001)		
Cd in urine (µg/L)	2003/04	201	0.60	0.15	2.15	0.48	0.44	0.34
	2004/05	189	0.70	0.27	1.91	(<0.0001)		
Creatinine in morning urine (g/L)	2003/04	201	0.72	0.15	1.80	-0.06	-0.47	-0.56
	2004/05	189	0.75	0.10	1.95	(0.40)		
Pb in blood (µg/L)	2003/04	201	90.90	47.10	213.00	0.70	0.69	0.62
	2004/05	189	85.00	39.10	203.00	(<0.0001)		
Se in serum (µg/L)	2003/04	201	56.90	36.00	78.50	0.45	0.17	0.05
	2004/05	189	72.10	46.40	95.70	(<0.0001)		

The intraclass correlation coefficient (ICC) is the between- subject minus the within- subject variance as a ratio to the sum of the two variances. A positive value of, for instance, 0.60 indicates that 60% of the variation is due to between-subject and 40% due to within-subject variance. ICC can become negative when the within-subject variance exceeds the between-subject variance.

Table 3. Clinical markers of kidney function in two consecutive investigations

Variable		N	Median	5%-value	95%-value	Spearman correlation coefficient (p)	ICC	ICC 5%-value																																																																																																												
Total protein in urine (mg/L)	2003/04	201	30.0	10	140	-0.03 (p=0.73)	0.03	-0.09																																																																																																												
	2004/05	189	13.5	0	86				Albumin in urine (mg/L)	2003/04	201	5.9	0.9	36	0.05 (0.53)	-0.14	-0.26	2004/05	188	1.0	0.9	17	β_2 -microglobulin in urine (ng/mL)	2003/04	201	25.1	7.3	171	0.41 (<0.0001)	-0.86	-0.89	2004/05	189	37.5	12.5	244	Creatinine in 1-hour urine (g/L)	2003/04	188	0.27	0.06	1.23	0.29 (p<0.001)	0.23	0.11	2004/05	189	0.44	0.11	1.7	Creatinine in 4-hour urine (g/l)	2003/04	201	0.19	0.05	1.43	0.05 (0.47)	-0.04	-0.16	2004/05	189	0.13	0.03	0.54	Glomerular filtration rate (4 h, measured)	2003/04	200	59.8	22.4	292	0.07 (0.33)	-0.16	-0.27	2004/05	189	41.5	5.6	144	CCR (Cockcroft and Gault) (mL/min)	2003/04	199	106.5	71.7	157	0.46 (<0.0001)	-0.3	-0.4	2004/05	189	99.2	62	187	Kidney length (average of both) (mm)	2003/04	201	118.5	106	128.5	0.91 (<0.0001)	0.92	0.9	2004/05	189	118	106.5	129.5	Minimal kidney cortex width (average) (mm)	2003/04	201	16.0	12	18.5	0.68 (<0.0001)	0.66	0.58	2004/05
Albumin in urine (mg/L)	2003/04	201	5.9	0.9	36	0.05 (0.53)	-0.14	-0.26																																																																																																												
	2004/05	188	1.0	0.9	17				β_2 -microglobulin in urine (ng/mL)	2003/04	201	25.1	7.3	171	0.41 (<0.0001)	-0.86	-0.89	2004/05	189	37.5	12.5	244	Creatinine in 1-hour urine (g/L)	2003/04	188	0.27	0.06	1.23	0.29 (p<0.001)	0.23	0.11	2004/05	189	0.44	0.11	1.7	Creatinine in 4-hour urine (g/l)	2003/04	201	0.19	0.05	1.43	0.05 (0.47)	-0.04	-0.16	2004/05	189	0.13	0.03	0.54	Glomerular filtration rate (4 h, measured)	2003/04	200	59.8	22.4	292	0.07 (0.33)	-0.16	-0.27	2004/05	189	41.5	5.6	144	CCR (Cockcroft and Gault) (mL/min)	2003/04	199	106.5	71.7	157	0.46 (<0.0001)	-0.3	-0.4	2004/05	189	99.2	62	187	Kidney length (average of both) (mm)	2003/04	201	118.5	106	128.5	0.91 (<0.0001)	0.92	0.9	2004/05	189	118	106.5	129.5	Minimal kidney cortex width (average) (mm)	2003/04	201	16.0	12	18.5	0.68 (<0.0001)	0.66	0.58	2004/05	189	15.5	12	19										
β_2 -microglobulin in urine (ng/mL)	2003/04	201	25.1	7.3	171	0.41 (<0.0001)	-0.86	-0.89																																																																																																												
	2004/05	189	37.5	12.5	244				Creatinine in 1-hour urine (g/L)	2003/04	188	0.27	0.06	1.23	0.29 (p<0.001)	0.23	0.11	2004/05	189	0.44	0.11	1.7	Creatinine in 4-hour urine (g/l)	2003/04	201	0.19	0.05	1.43	0.05 (0.47)	-0.04	-0.16	2004/05	189	0.13	0.03	0.54	Glomerular filtration rate (4 h, measured)	2003/04	200	59.8	22.4	292	0.07 (0.33)	-0.16	-0.27	2004/05	189	41.5	5.6	144	CCR (Cockcroft and Gault) (mL/min)	2003/04	199	106.5	71.7	157	0.46 (<0.0001)	-0.3	-0.4	2004/05	189	99.2	62	187	Kidney length (average of both) (mm)	2003/04	201	118.5	106	128.5	0.91 (<0.0001)	0.92	0.9	2004/05	189	118	106.5	129.5	Minimal kidney cortex width (average) (mm)	2003/04	201	16.0	12	18.5	0.68 (<0.0001)	0.66	0.58	2004/05	189	15.5	12	19																								
Creatinine in 1-hour urine (g/L)	2003/04	188	0.27	0.06	1.23	0.29 (p<0.001)	0.23	0.11																																																																																																												
	2004/05	189	0.44	0.11	1.7				Creatinine in 4-hour urine (g/l)	2003/04	201	0.19	0.05	1.43	0.05 (0.47)	-0.04	-0.16	2004/05	189	0.13	0.03	0.54	Glomerular filtration rate (4 h, measured)	2003/04	200	59.8	22.4	292	0.07 (0.33)	-0.16	-0.27	2004/05	189	41.5	5.6	144	CCR (Cockcroft and Gault) (mL/min)	2003/04	199	106.5	71.7	157	0.46 (<0.0001)	-0.3	-0.4	2004/05	189	99.2	62	187	Kidney length (average of both) (mm)	2003/04	201	118.5	106	128.5	0.91 (<0.0001)	0.92	0.9	2004/05	189	118	106.5	129.5	Minimal kidney cortex width (average) (mm)	2003/04	201	16.0	12	18.5	0.68 (<0.0001)	0.66	0.58	2004/05	189	15.5	12	19																																						
Creatinine in 4-hour urine (g/l)	2003/04	201	0.19	0.05	1.43	0.05 (0.47)	-0.04	-0.16																																																																																																												
	2004/05	189	0.13	0.03	0.54				Glomerular filtration rate (4 h, measured)	2003/04	200	59.8	22.4	292	0.07 (0.33)	-0.16	-0.27	2004/05	189	41.5	5.6	144	CCR (Cockcroft and Gault) (mL/min)	2003/04	199	106.5	71.7	157	0.46 (<0.0001)	-0.3	-0.4	2004/05	189	99.2	62	187	Kidney length (average of both) (mm)	2003/04	201	118.5	106	128.5	0.91 (<0.0001)	0.92	0.9	2004/05	189	118	106.5	129.5	Minimal kidney cortex width (average) (mm)	2003/04	201	16.0	12	18.5	0.68 (<0.0001)	0.66	0.58	2004/05	189	15.5	12	19																																																				
Glomerular filtration rate (4 h, measured)	2003/04	200	59.8	22.4	292	0.07 (0.33)	-0.16	-0.27																																																																																																												
	2004/05	189	41.5	5.6	144				CCR (Cockcroft and Gault) (mL/min)	2003/04	199	106.5	71.7	157	0.46 (<0.0001)	-0.3	-0.4	2004/05	189	99.2	62	187	Kidney length (average of both) (mm)	2003/04	201	118.5	106	128.5	0.91 (<0.0001)	0.92	0.9	2004/05	189	118	106.5	129.5	Minimal kidney cortex width (average) (mm)	2003/04	201	16.0	12	18.5	0.68 (<0.0001)	0.66	0.58	2004/05	189	15.5	12	19																																																																		
CCR (Cockcroft and Gault) (mL/min)	2003/04	199	106.5	71.7	157	0.46 (<0.0001)	-0.3	-0.4																																																																																																												
	2004/05	189	99.2	62	187				Kidney length (average of both) (mm)	2003/04	201	118.5	106	128.5	0.91 (<0.0001)	0.92	0.9	2004/05	189	118	106.5	129.5	Minimal kidney cortex width (average) (mm)	2003/04	201	16.0	12	18.5	0.68 (<0.0001)	0.66	0.58	2004/05	189	15.5	12	19																																																																																
Kidney length (average of both) (mm)	2003/04	201	118.5	106	128.5	0.91 (<0.0001)	0.92	0.9																																																																																																												
	2004/05	189	118	106.5	129.5				Minimal kidney cortex width (average) (mm)	2003/04	201	16.0	12	18.5	0.68 (<0.0001)	0.66	0.58	2004/05	189	15.5	12	19																																																																																														
Minimal kidney cortex width (average) (mm)	2003/04	201	16.0	12	18.5	0.68 (<0.0001)	0.66	0.58																																																																																																												
	2004/05	189	15.5	12	19																																																																																																															

Table 4. Association of metals and metalloids with β_2 -microglobulin, creatinine clearance, and kidney length controlling for confounders

	β_2 -microglobulin			CCR Cockcroft, Gault			Kidney length (mm)		
	Total sample n=201, obs=387	Ben- offspring n=102, obs=192	Control- offspring n=99, obs=195	Total sample n=201, obs=388	Ben- offspring n=102, obs=193	Control- offspring n=99, obs=195	Total sample n=201, obs=388	Ben- offspring n=102, obs=193	Control- offspring n=99, obs=195
Intercept	1.260	0.992	1.358	121.00	98.97	134.12	100.32	98.41	99.91
Blood lead ($\mu\text{g/L}$)	-0.001	-0.001	-0.001	-0.03	0.02	-0.09*	0.005	-0.01	0.00
Selenium in serum ($\mu\text{g/L}$)	0.001	0.003*	0.000	-0.21*	-0.15	-0.29*	0.02*	0.04*	0.01
Cadmium in urine ($\mu\text{g/L}$)	0.042	0.074	0.003	-2.19	-2.64	-2.68	0.10	0.11	0.05
Arsenic in urine ($\mu\text{g/L}$)	0.001	0.001	0.002	0.25	0.82	0.18	-0.06	-0.11	-0.07
Sex, males	0.189*	0.235*	0.151*	8.52*	0.54	16.21*	1.92	1.19	2.45
Age (years)	0.004*	0.008*	0.002	-1.12*	-0.89*	-1.24*	0.01	-0.02	0.02
Current smoker	-0.095*	-0.233*	-0.004	-0.56	3.37	-3.76	0.29	2.77*	-0.48
Ex-smoker	0.024	-0.059	0.039	-3.92	4.20	-14.02*	-0.12	0.98	-0.09
Diabetes mellitus	0.058	0.210	-0.063	-4.93	-7.87	-7.10	-1.87	-4.01	-0.25
Hypertension	0.048	-0.040	0.100	-	-	-	-0.89	0.53	-1.72
Body surface area #	-0.174	-0.139	-0.116	30.41*	31.99*	31.58*	8.58*	8.74*	9.55*
Cystitis	0.014	0.032	-0.010	-1.68	3.59	-3.04	-0.67	-1.07	-0.42
Pyelonephritis	-0.081	-0.118	0.191	3.66	3.08	-23.74	-0.65	0.77	-0.50
Kidney stone(s)	0.017	0.014	0.123	4.26	4.71	4.29	-0.57	-1.53	0.18
Creatinine in 1-hour urine	0.475*	0.417*	0.530*	0.001*	0.002*	0.001	-	-	-
Creatinine in morning urine (g/L)	0.018	0.004	0.006	-3.71	-2.25	-5.19	-0.07	0.35	-0.28

* $p \leq 0.05$

BSA = square root ((height in cm \times weight in kg)/3600)